

## AN ALGEBRAIC MODEL FOR CHAINS ON $\Omega BG_p^\wedge$

DAVE BENSON

ABSTRACT. We provide an interpretation of the homology of the loop space on the  $p$ -completion of the classifying space of a finite group in terms of representation theory, and demonstrate how to compute it. We then give the following reformulation. If  $f$  is an idempotent in  $kG$  such that  $f.kG$  is the projective cover of the trivial module  $k$ , and  $e = 1 - f$ , then we exhibit isomorphisms for  $n \geq 2$ :

$$\begin{aligned} H_n(\Omega BG_p^\wedge; k) &\cong \mathrm{Tor}_{n-1}^{e.kG, e}(kG.e, e.kG), \\ H^n(\Omega BG_p^\wedge; k) &\cong \mathrm{Ext}_{e.kG, e}^{n-1}(e.kG, e.kG). \end{aligned}$$

Further algebraic structure is examined, such as products and coproducts, restriction and Steenrod operations.

### 1. INTRODUCTION

This paper grew out of an attempt to understand the relationship between the work of Ran Levi [17, 18, 19, 20] (see also Cohen and Levi [6]) on  $H_*(\Omega BG_p^\wedge; k)$ , the homology of the loop space on the Bousfield–Kan  $p$ -completion of the classifying space of a finite group  $G$  with coefficients in a field  $k$ , and the papers [8, 9] of Dwyer, Greenlees and Iyengar. The result is a representation theoretic description of the homology of  $\Omega BG_p^\wedge$ . For reasonably small groups, this is explicitly computable, and to illustrate the methods we give the results of some computations using the computer algebra system MAGMA [4].

In order to describe our results, we introduce the following operation in modular representation theory. Let  $k$  be a field of characteristic  $p$ , and let  $G$  be a finite group. Recall that  $O^p(G)$  is the smallest normal subgroup of  $G$  such that the quotient is a  $p$ -group. If  $M$  is a finitely generated  $kG$ -module, we define  $[O^p(G), M]$  to be the  $k$ -linear span of the set

$$\{g(m) - m \mid g \in O^p(G), m \in M\}.$$

This is the smallest submodule of  $M$  such that  $O^p(G)$  acts trivially on the quotient, or equivalently the smallest submodule such that the quotient has a filtration where  $G$  acts trivially on the filtered quotients (see Lemma 3.1).

Inductively define  $P_0 = N_0$  to be the projective cover  $P(k)$  of the trivial  $kG$ -module  $k$ , and for  $i \geq 1$ ,  $M_{i-1} = [O^p(G), N_{i-1}]$ ,  $P_i$  is the projective cover of  $M_{i-1}$ , and  $N_i = \Omega(M_{i-1})$ , the kernel of  $P_i \rightarrow M_{i-1}$ . This construction gives us a complex

---

Received by the editors July 25, 2007.

2000 *Mathematics Subject Classification*. Primary 55P35, 55R35, 20C20; Secondary 55P60, 20J06, 13C40, 14M10.

©2008 American Mathematical Society  
 Reverts to public domain 28 years from publication

of projective  $kG$ -modules

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow 0$$

with the property that  $O^p(G)$  acts trivially on the homology, and the  $P_i$  with  $i \geq 1$  do not have the projective cover of  $k$  as a direct summand. We call this a *left  $k$ -squeezed resolution* for  $G$ , and we write  $H_*^\Omega(G, k)$  for the homology of this complex. A more formal definition is given in Section 3.

The dual of this construction gives us a complex of injective  $kG$ -modules

$$0 \rightarrow I_0 \rightarrow I_{-1} \rightarrow I_{-2} \rightarrow \cdots$$

such that  $I_0$  is the injective hull of the trivial module (which is isomorphic to  $P(k)$ ),  $O^p(G)$  acts trivially on the homology, and the  $I_{-i}$  with  $i \geq 1$  do not have  $P(k)$  as a direct summand. We call this a *right  $k$ -squeezed resolution* for  $G$ , and we write  $H_\Omega^*(G, k)$  for the homology of this complex, with the indexing negated to make it cohomological.

Our first theorem, which we prove in Section 5, gives a homological interpretation for the construction. Let  $f$  be an idempotent in  $kG$  such that  $f.kG$  is isomorphic to the projective cover of the trivial module and set  $e = 1 - f$ .

**Theorem 1.1.** *For  $n \geq 2$ , we have isomorphisms*

$$\begin{aligned} \text{Tor}_{n-1}^{e.kG.e}(kG.e, e.kG) &\cong H_n^\Omega(G, k), \\ \text{Ext}_{e.kG.e}^{n-1}(e.kG, e.kG) &\cong H_\Omega^n(G, k). \end{aligned}$$

*There are exact sequences*

$$\begin{aligned} 0 \rightarrow H_1^\Omega(G, k) \rightarrow kG.e \otimes_{e.kG.e} e.kG \rightarrow kG \rightarrow H_0^\Omega(G, k) \rightarrow 0, \\ 0 \rightarrow H_\Omega^0(G, k) \rightarrow kG \rightarrow \text{Hom}_{e.kG.e}(e.kG, e.kG) \rightarrow H_\Omega^1(G, k) \rightarrow 0. \end{aligned}$$

Notice that we can rewrite the left hand sides of the isomorphisms in Theorem 1.1 as

$$\text{Tor}_{n-1}^{e.kG.e}(f.kG.e, e.kG.f) \quad \text{and} \quad \text{Ext}_{e.kG.e}^{n-1}(e.kG.f, e.kG.f).$$

This makes computations slightly easier because  $e.kG.f$  and  $f.kG.e$  are smaller than  $e.kG$  and  $kG.e$ .

Our second theorem gives a topological interpretation of  $H_*^\Omega(G, k)$  and  $H_\Omega^*(G, k)$ .

**Theorem 1.2.** *For  $i \geq 0$ ,  $H_i^\Omega(G, k) \cong H_i(\Omega BG_p^\wedge; k)$  and  $H_\Omega^i(G, k) \cong H^i(\Omega BG_p^\wedge; k)$ .*

It is this theorem that we use for the MAGMA computations; examples can be found in the next section.

We prove Theorem 1.2 in Section 6 by making a model of  $\Omega BG_p^\wedge$  that admits a free action of  $G$  with acyclic quotient. In Section 7, we provide another proof by interpreting the chains on  $\Omega BG_p^\wedge$  as a best approximation to  $kG$  by chain complexes built from an injective resolution of  $k$ . This proof uses more machinery, but makes it easy to interpret the loop multiplication in  $H_*(\Omega BG_p^\wedge; k)$  algebraically.

Combining the statements of Theorems 1.1 and 1.2, we obtain isomorphisms for  $n \geq 2$ :

$$(1.3) \quad \text{Tor}_{n-1}^{e.kG.e}(f.kG.e, e.kG.f) \cong H_n(\Omega BG_p^\wedge; k),$$

$$(1.4) \quad \text{Ext}_{e.kG.e}^{n-1}(e.kG.f, e.kG.f) \cong H^n(\Omega BG_p^\wedge; k).$$

Finally, in the last section we discuss some of the problems that motivated this work. These involve the boundary between polynomial and exponential growth for

$H_*(\Omega BG_p^\wedge; k)$ . We completely solve the problem in characteristic two for groups with an elementary abelian Sylow 2-subgroup  $E$ . In this situation, we show that polynomial growth occurs if and only if  $H^*(G, k)$  is a complete intersection. This condition is equivalent to the statement that  $N_G(E)/O_2'N_G(E)$  is a direct product of copies of  $\mathbb{Z}/2$ ,  $A_4$  and  $(\mathbb{Z}/2)^3 \rtimes F_{21}$ , where  $F_{21}$  is a Frobenius group of order 21.

2. EXAMPLES

We illustrate the main theorems with some examples.

**Example 2.1.** Let  $G = A_4$ , the alternating group of degree four, and let  $k$  be a field of characteristic two. We write  $k, \omega$  and  $\bar{\omega}$  for the three simple  $kG$ -modules. The projective indecomposable  $kG$ -modules have the following structure:

$$\begin{array}{ccccccc}
 & k & & \omega & & \bar{\omega} & \\
 \omega & & \bar{\omega} & & k & & \omega \\
 & k & & \omega & & \bar{\omega} & \\
 P(k) & & & P(\omega) & & & P(\bar{\omega})
 \end{array}$$

We have  $P_0 = N_0 = P(k)$ ,

$$\begin{array}{l}
 M_0 = \begin{array}{ccc} \omega & & \bar{\omega} \\ & k & \end{array} \quad P_1 = P(\omega) \oplus P(\bar{\omega}) \quad N_1 = \begin{array}{ccc} \bar{\omega} & & k \\ & \omega & \bar{\omega} \end{array} \omega \\
 M_1 = \begin{array}{ccc} \bar{\omega} & & \omega \\ & \omega & \bar{\omega} \end{array} \oplus \begin{array}{ccc} \omega & & \bar{\omega} \\ & \bar{\omega} & \omega \end{array} \quad P_2 = P(\bar{\omega}) \oplus P(\omega) \quad N_2 = \begin{array}{ccc} k & & k \\ & \bar{\omega} & \omega \end{array} \oplus \begin{array}{ccc} k & & \omega \\ & \omega & \bar{\omega} \end{array} \\
 M_2 = \bar{\omega} \oplus \omega \quad P_3 = P(\bar{\omega}) \oplus P(\omega) \quad N_3 = \begin{array}{ccc} \bar{\omega} & & k \\ & \omega & \bar{\omega} \end{array} \oplus \begin{array}{ccc} k & & \omega \\ & \bar{\omega} & \omega \end{array}
 \end{array}$$

Thereafter, the construction has period two, so that  $M_i \cong M_{i-2}$ ,  $P_i \cong P_{i-2}$  and  $N_i \cong N_{i-2}$ . So according to Theorem 1.2,  $H_n(\Omega BG_p^\wedge; k)$  has dimension one for  $n = 0$  and  $n = 1$ , and it has dimension two if  $n \geq 2$ .

It is shown in Levi [19] that  $H_*(\Omega BG_p^\wedge; k)$  does not always have polynomial growth. Using the computer system MAGMA [4], some examples of exponential growth were investigated using Theorem 1.2, and the following table gives the dimensions of the first few homology groups for these examples.

		Degree										
Group	$p$	0	1	2	3	4	5	6	7	8	9	10
$(\mathbb{Z}/2)^3 \rtimes \mathbb{Z}/7$	2	1	0	1	3	3	6	12	18	33	57	96
$(\mathbb{Z}/3)^2 \rtimes \mathbb{Z}/2$	3	1	1	5	12	32	84	220	576			
$(\mathbb{Z}/3)^2 \rtimes \mathbb{Z}/4$	3	1	1	3	6	12	22	42	80	150		
$(\mathbb{Z}/5)^2 \rtimes \mathbb{Z}/3$	5	1	1	3	6	14	30	64	138			

In each case the action in the semidirect product is faithful, and in the second entry the involution acts to invert every element of  $(\mathbb{Z}/3)^2$ .

**Example 2.2.** Take the group  $(\mathbb{Z}/3)^2 \rtimes \mathbb{Z}/2$  given by the second entry in the table above, with  $k$  a field of characteristic three. Then  $A = e.kG.e$  is the five dimensional commutative algebra

$$A = k[u, v, w]/(uw - v^2, u^2, uv, vw, w^2).$$


This is a Koszul algebra, whose Koszul dual is

$$A^1 = \text{Ext}_A^*(k, k) = k\langle \alpha, \beta, \gamma \rangle / (\alpha\gamma + \gamma\alpha + \beta^2).$$

The module  $M = e.kG.f$  is the four dimensional module given in terms of generators and relations as

$$M = (Ax \oplus Ay) / (vx - uy, wx - vy, ux, wy).$$

Here are pictures of  $A$  and  $M$ :



It is not hard to compute the syzygies of  $M$  as an  $A$ -module. Since  $A$  is a symmetric algebra with  $J(A)^3 = 0$ , each  $\Omega^n M$  is an indecomposable module with  $\text{Rad}^2 \Omega^n M = 0$  and  $\text{Soc}(\Omega^n M) = \text{Rad}(\Omega^n M) \cong \Omega^{n-1} M / \text{Rad}(\Omega^{n-1} M)$ . Set  $b_n = \dim_k \Omega^n M / \text{Rad}(\Omega^n M)$  and  $c_n = \dim_k \text{Rad}(\Omega^n M)$ . Then we have  $b_0 = c_0 = 2$ ,  $c_{n+1} = b_n$  and  $b_{n+1} = 3b_n - c_n$ .

This gives us the recurrence relation

$$b_0 = 2, \quad b_1 = 4, \quad b_{n+1} = 3b_n - b_{n-1} \quad (n \geq 2),$$

and hence the generating function

$$\sum_{n \geq 0} b_n t^n = \frac{2(1-t)}{1-3t+t^2} = 2 + 4t + 10t^2 + 26t^3 + 68t^4 + \dots$$

Thus  $b_n = 2F_{2n+1}$ , where  $F_n$  is the  $n$ th Fibonacci number.

**Lemma 2.3.** For every homomorphism from  $\Omega^n M$  to  $M$ , for  $n \geq 1$ , the image lies in  $\text{Rad}(M) = \text{Soc}(M)$ .

*Proof (outline).* The first step is to prove that there are short exact sequences

$$0 \rightarrow \Omega^n M \rightarrow \Omega^{n+1} M \rightarrow \Omega^n k \oplus \Omega^n k \rightarrow 0$$

for all  $n \geq 0$ . The second step is to prove by induction on  $n$  that for every  $n \geq 0$ , every endomorphism of  $\Omega^n M$  can be written as a multiple of the identity plus an endomorphism whose image lies in  $\text{Rad}(\Omega^n M)$ . The final step is to compose the given homomorphism from  $\Omega^n M$  to  $M$  with the composite of the inclusions given by the short exact sequences above,  $M \hookrightarrow \Omega M \hookrightarrow \dots \hookrightarrow \Omega^n M$  to obtain an endomorphism of  $\Omega^n M$ , and examine the image. □

It follows from the lemma that for  $n \geq 2$  there are short exact sequences

$$0 \rightarrow \text{Ext}_A^{n-1}(M, M/\text{Rad}(M)) \rightarrow \text{Ext}_A^n(M, \text{Rad}(M)) \rightarrow \text{Ext}_A^n(M, M) \rightarrow 0.$$

Thus for  $n \geq 2$  we have  $\dim_k \text{Ext}_A^n(M, M) = 2b_n - 2b_{n-1} = 4F_{2n}$ . So

$$\sum_{n \geq 2} \dim_k \text{Ext}_A^n(M, M) = -2b_1 t^2 + 2(1-t) \sum_{n \geq 2} b_n t^n.$$

Finally, adjusting to give the correct dimensions for  $n \leq 2$ , Theorem 1.1 gives us

$$\begin{aligned} \sum_{n \geq 0} \dim_k H^n(\Omega BG_p^\wedge; k)t^n &= 1 + t + 5t^2 + 2t(1 - t) \sum_{n \geq 2} b_n t^n \\ &= \frac{(1 - t + t^2)^2}{1 - 3t + t^2} \\ &= 1 + t + 5t^2 + 12t^3 + 32t^4 + 84t^5 + 220t^6 + \dots \end{aligned}$$

This function has poles at  $(3 \pm \sqrt{5})/2$ ; since one of these is in the interior of the unit circle in the complex plane, it follows that  $\dim_k H^n(\Omega BG_p^\wedge; k)$  is growing exponentially with  $n$ . Alternatively, note that for  $n \geq 3$  we have  $\dim_k H^n(\Omega BG_p^\wedge; k) = 4F_{2n-2}$ , and the Fibonacci numbers grow exponentially.

### 3. SQUEEZED RESOLUTIONS

Let  $G$  be a finite group, and  $k$  be a field of characteristic  $p$ . We also denote by  $k$  the trivial  $kG$ -module; namely the ring  $k$  regarded as a  $kG$ -module with trivial  $G$ -action. Thus if  $I_G$  is the augmentation ideal of  $kG$ , generated by the elements  $g - 1$  for  $g \in G$ , then  $k \cong kG/I_G$ .

If  $M$  is a  $kG$ -module, we define

$$[O^p(G), M] = \{g(m) - m \mid g \in O^p(G), m \in M\}.$$

**Lemma 3.1.**  *$[O^p(G), M]$  is the smallest submodule of  $M$  such that the quotient has a filtration with  $G$  acting trivially on the filtered quotients.*

*Proof.*  $O^p(G)$  acts trivially on  $M/[O^p(G), M]$ , and  $G/O^p(G)$  is a  $p$ -group. So the filtration whose  $i$ th term is  $J^i(G/O^p(G))(M/[O^p(G), M])$  has  $G$  acting trivially on the filtered quotients. Conversely, suppose that  $M'$  is a submodule of  $M$  such that  $M/M'$  has a filtration

$$M/M' = M_0 \supseteq M_1 \supseteq \dots \supseteq M_n = \{0\}$$

with  $G$  acting trivially on  $M_i/M_{i+1}$  ( $0 \leq i \leq n - 1$ ). Since  $O^p(G)$  admits no homomorphisms to  $k^+$ ,  $\text{Ext}_{kO^p(G)}^1$  vanishes on pairs of modules with trivial  $O^p(G)$ -action. So the filtration splits on restriction to  $O^p(G)$ . Hence  $O^p(G)$  acts trivially on  $M/M'$ , and so  $M' \supseteq [O^p(G), M]$ .  $\square$

We have

$$\text{Hom}_{kG}([O^p(G), M], k) = 0.$$

Dually, we define  $M^{O^p(G)}$  to be the fixed points of  $O^p(G)$  on  $M$ . This is the largest submodule of  $M$  that has a filtration such that  $G$  acts trivially on the filtered quotients. We have

$$\text{Hom}_{kG}(k, M/M^{O^p(G)}) = 0.$$

**Definition 3.2.** A *left  $k$ -squeezed resolution* is a complex

$$\dots \rightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \rightarrow 0$$

of  $kG$ -modules satisfying the following conditions:

- (i) each  $P_n$  is projective,
- (ii)  $H_n(P_* \otimes_{kG} k) \cong \begin{cases} k, & n = 0, \\ 0, & n \neq 0, \end{cases}$
- (iii)  $[O^p(G), H_n(P_*)] = 0$  for all  $n$ .

We can construct a complex satisfying these conditions inductively by the process described in the Introduction. The dual definition is as follows.

**Definition 3.3.** A *right  $k$ -squeezed resolution* is a complex

$$0 \rightarrow I_0 \xrightarrow{d_0} I_{-1} \xrightarrow{d_{-1}} I_{-2} \xrightarrow{d_{-2}} \dots$$

of  $kG$ -modules satisfying the following conditions:

- (i) each  $I_n$  is injective,
- (ii)  $H_n \operatorname{Hom}_{kG}(k, I_*) \cong \begin{cases} k, & n = 0, \\ 0, & n \neq 0, \end{cases}$
- (iii)  $[O^p(G), H_n(I_*)] = 0$  for all  $n$ .

We can construct a complex satisfying these conditions inductively by the dual of the process described in the Introduction.

**Theorem 3.4** (Comparison Theorem). (i) *Let*

$$\begin{aligned} \dots &\rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow 0, \\ \dots &\rightarrow P'_2 \rightarrow P'_1 \rightarrow P'_0 \rightarrow 0 \end{aligned}$$

*be complexes together with augmentations  $P_0 \rightarrow k$  and  $P'_0 \rightarrow k$  satisfying:*

- (a) *each  $P_i$  is a projective  $kG$ -module, and the augmentation  $P_0 \rightarrow k$  induces an isomorphism  $H_*(P_* \otimes_{kG} k) \rightarrow k$ ;*
- (b)  $[O^pG, H_*(P'_*)] = 0$ .

*Then there exists a map of complexes  $P_* \rightarrow P'_*$  lifting the identity map on  $k$ :*

$$\begin{array}{ccccccccc} \dots & \longrightarrow & P_2 & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & k & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \\ \dots & \longrightarrow & P'_2 & \longrightarrow & P'_1 & \longrightarrow & P'_0 & \longrightarrow & k & \longrightarrow & 0 \end{array}$$

*Any two such comparison maps are chain homotopic.*

(ii) *Let*

$$\begin{aligned} 0 &\rightarrow I_0 \rightarrow I_{-1} \rightarrow I_{-2} \rightarrow \dots, \\ 0 &\rightarrow I'_0 \rightarrow I'_{-1} \rightarrow I'_{-2} \rightarrow \dots \end{aligned}$$

*be complexes together with coaugmentations  $k \rightarrow I_0$  and  $k \rightarrow I'_0$  satisfying:*

- (a)  $[O^pG, H_*(I_*)] = 0$ ;
- (b) *each  $I'_{-i}$  is an injective  $kG$ -module, and the augmentation  $k \rightarrow I'_0$  induces an isomorphism  $k \rightarrow H_* \operatorname{Hom}_{kG}(k, I'_*)$ .*

*Then there exists a map of complexes  $I_* \rightarrow I'_*$  extending the identity map on  $k$ :*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & k & \longrightarrow & I_0 & \longrightarrow & I_{-1} & \longrightarrow & I_{-2} & \longrightarrow & \dots \\ & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & k & \longrightarrow & I'_0 & \longrightarrow & I'_{-1} & \longrightarrow & I'_{-2} & \longrightarrow & \dots \end{array}$$

*Any two such comparison maps are chain homotopic.*

*Proof.* This is an exercise in diagram chasing. □

**Corollary 3.5.** (i) *Given two left  $k$ -squeezed resolutions  $P_*$  and  $P'_*$ , an isomorphism*

$$\mathrm{Hom}_{kG}(P_0, k) \cong \mathrm{Hom}_{kG}(P'_0, k)$$

*lifts to a chain homotopy equivalence  $P_* \rightarrow P'_*$ , and hence to an isomorphism*

$$H_*(P_*) \cong H_*(P'_*).$$

(ii) *Given two right  $k$ -squeezed resolutions  $I_*$  and  $I'_*$ , an isomorphism*

$$\mathrm{Hom}_{kG}(k, I_0) \cong \mathrm{Hom}_{kG}(k, I'_0)$$

*extends to a chain homotopy equivalence  $I_* \rightarrow I'_*$ , and hence to an isomorphism*

$$H_*(I_*) \cong H_*(I'_*).$$

*Proof.* This follows from the theorem. □

**Definition 3.6.** Given a finite group  $G$ , Corollary 3.5 implies that the homology of a left  $k$ -squeezed resolution  $P_*$  does not depend on the choice of resolution. So we define  $H_n^\Omega(G, k) = H_n(P_*)$ . Similarly, if  $I_*$  is a right  $k$ -squeezed resolution, we define  $H_n^\Omega(G, k) = H_{-n}(I_*)$ .

#### 4. PRODUCTS, COPRODUCTS, SUBGROUPS, STEENROD OPERATIONS

**4.1. External coproducts.** Let  $G$  and  $G'$  be a finite group. If  $P_*$  and  $P'_*$  are left  $k$ -squeezed resolutions for  $G$  and  $G'$ , then  $P_* \otimes_k P'_*$  is a left  $k$ -squeezed resolution for  $G \times G'$ . This gives a Künneth theorem for homology. This gives

$$H_*^\Omega(G, k) \otimes_k H_*^\Omega(G', k) \cong H_*^\Omega(G \times G', k).$$

Dually, if  $I_*$  and  $I'_*$  are right  $k$ -squeezed resolutions for  $G$  and  $G'$ , then  $I_* \otimes_k I'_*$  is a right  $k$ -squeezed resolution for  $G \times G'$ . This gives

$$H_\Omega^*(G, k) \otimes_k H_\Omega^*(G', k) \cong H_\Omega^*(G \times G', k).$$

**4.2. Restriction and corestriction maps.** Let  $H$  be a subgroup of a finite group  $G$ . If  $P_*$  is a left  $k$ -squeezed resolution for  $G$  and  $Q_*$  is a left  $k$ -squeezed resolution for  $H$  then the Comparison Theorem, Theorem 3.4, gives a map  $Q_* \rightarrow \mathrm{res}_{G,H} P_*$  and hence a corestriction map

$$\mathrm{cores}_{H,G}: H_*^\Omega(H, k) \rightarrow H_*^\Omega(G, k).$$

Dually, there is a restriction map

$$\mathrm{res}_{G,H}: H_\Omega^*(G, k) \rightarrow H_\Omega^*(H, k).$$

**4.3. Internal coproducts.** Internal coproducts can be obtained by taking the external product for  $G \times G$  and then restricting to the diagonal copy of  $G$ . This gives rise to a comparison map  $P_* \rightarrow P_* \otimes P_*$  and hence to cocommutative coproducts

$$(4.4) \quad H_*^\Omega(G, k) \rightarrow H_*^\Omega(G, k) \otimes_k H_*^\Omega(G, k).$$

Dually, there are graded commutative products

$$(4.5) \quad H_\Omega^*(G, k) \otimes_k H_\Omega^*(G, k) \rightarrow H_\Omega^*(G, k).$$

**4.6. Products.** Using the Comparison Theorem, Theorem 3.4, an element of  $H_n^\Omega(G, k)$  can be lifted to a map of complexes

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & P_{n+1} & \longrightarrow & P_n & \longrightarrow & P_{n-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow 0 \end{array}$$

which is unique up to chain homotopy. This allows us to use composition of maps to define a product

$$(4.7) \quad H_*^\Omega(G, k) \otimes_k H_*^\Omega(G, k) \rightarrow H_*^\Omega(G, k).$$

This product is associative, but usually not graded commutative. In Section 7 we shall see that this product corresponds to the product on  $H_*(\Omega BG_p^\wedge; k)$  coming from the composition of loops.

**4.8. Steenrod operations.** The comparison map (4.4) is  $E_\infty$ . In particular, if  $k$  is a field, then there is an action of the Steenrod algebra on  $H_*^\Omega(G, k)$  lowering degree. Similarly, there is an action of the Steenrod algebra on  $H_\Omega^*(G, k)$  raising degree. We may develop this theory algebraically by analogy with the Evens norm map [1, Chapter 4] as follows.

Let  $k$  be a field of characteristic  $p$ . Suppose that  $G$  is a finite group, and consider the group  $G \times \mathbb{Z}/p$ . Then we have  $O^p(G \times \mathbb{Z}/p) = O^p(G)$ . So if  $J_*$  is a right  $k$ -squeezed resolution for  $G$ , then the tensor induced complex  $J_* \otimes^{G \times \mathbb{Z}/p}$  satisfies  $[O^p(G), H_*(J_* \otimes^{G \times \mathbb{Z}/p})] = 0$ . There is a subtle point here coming from the signs. If  $p = 2$  the signs are not a problem, but if  $p$  is odd, then we must remark that the action of  $G \times \mathbb{Z}/p$  on the cosets of  $G$  lies in the alternating group, and so the action of  $G$  on the homology of the tensor product is trivial.

It follows using the Comparison Theorem, Theorem 3.4, that if  $I_*$  is a right  $k$ -squeezed resolution for  $G$  and  $Z_*$  is an injective resolution of  $k$  for  $\mathbb{Z}/p$ , then there is a comparison map  $J_* \otimes^{G \times \mathbb{Z}/p} \rightarrow I_* \otimes_k Z_*$ . So if  $x \in H_\Omega^n(G, k) = H_{-n}(J_*)$ , then  $x^{\otimes p}$  defines an element of  $H_{-np}(J_* \otimes^{G \times \mathbb{Z}/p})$  and hence by composition an element of  $H_{-np}(I_* \otimes_k Z_*)$ . It may be checked as in Lemma 4.1.1 of [1] that the resulting element is independent of the representative of the homology class, and so we obtain a well defined map

$$\text{Norm}_{G, G \times \mathbb{Z}/p}: H_\Omega^n(G, k) \rightarrow H_\Omega^*(G, k) \otimes_k H^*(\mathbb{Z}/p, k).$$

As in Section 4.4 of [1], picking out coefficients of elements of  $H^*(\mathbb{Z}/p, k)$  gives the Steenrod operations on  $H_\Omega^*(G, k)$ . For example if  $k = \mathbb{F}_2$ , then we have  $H^*(\mathbb{Z}/2, \mathbb{F}_2) = \mathbb{F}_2[t]$  with  $|t| = 1$ . So if  $x \in H_\Omega^n(G, \mathbb{F}_2)$ , then

$$\text{Norm}_{G, G \times \mathbb{Z}/2}(x) = \sum_{j=0}^n \text{Sq}^j(x) \otimes t^{n-j}$$

with  $\text{Sq}^j(x) \in H^{n+j}(G, \mathbb{F}_2)$ . The corresponding formulae for  $p$  odd are more complicated, but are essentially the same as in [1, Proposition 4.4.3].

**4.9. Yoneda interpretation.** An  $n$ -fold  $k$ -squeezed extension for  $G$  is a complex of  $kG$ -modules

$$(4.10) \quad M_*: \quad 0 \rightarrow k \rightarrow M_{n-1} \rightarrow \cdots \rightarrow M_0 \rightarrow k \rightarrow 0$$

such that

- (i)  $[O^p(G), H_*(M_*)] = 0$ ,
- (ii) the map  $k \rightarrow M_{n-1}$  is injective, and
- (iii)  $M_0 \rightarrow k$  is surjective.

A map of  $n$ -fold  $k$ -squeezed extensions is a map of complexes which is the identity on the two copies of  $k$ . Two  $n$ -fold  $k$ -squeezed extensions are equivalent if there is a chain of maps connecting one with the other; in other words, this is the equivalence relation generated by the existence of a map of extensions.

Let  $P_*$  be a left  $k$ -squeezed resolution for  $G$ . Given an  $n$ -fold  $k$ -squeezed extension as above, the comparison theorem gives us a map of complexes

$$\begin{array}{ccccccc}
 P_{n-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \longrightarrow & k \longrightarrow 0 \\
 \downarrow & & & & \downarrow & & \downarrow \\
 0 & \longrightarrow & k & \longrightarrow & M_{n-1} & \longrightarrow & \cdots \longrightarrow M_0 \longrightarrow k \longrightarrow 0
 \end{array}$$

and hence a map  $\hat{\zeta}: \text{Ker}(P_{n-1} \rightarrow P_{n-2}) \rightarrow k$ . The map  $\hat{\zeta}$  gives us a well defined element  $\zeta \in H_\Omega^n(G, k)$  that is an invariant of the equivalence class of the  $k$ -squeezed extension. Writing  $L_\zeta$  for the kernel of  $\hat{\zeta}$ , we obtain a map of  $k$ -squeezed extensions

$$\begin{array}{ccccccc}
 0 & \longrightarrow & k & \longrightarrow & P_{n-1}/L_\zeta & \longrightarrow & \cdots \longrightarrow P_0 \longrightarrow k \longrightarrow 0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & k & \longrightarrow & M_{n-1} & \longrightarrow & \cdots \longrightarrow M_0 \longrightarrow k \longrightarrow 0
 \end{array}$$

This way, we see that the equivalence classes of  $n$ -fold  $k$ -squeezed extensions are in one-to-one correspondence with elements of  $H_\Omega^n(G, k)$ .

**4.11. The antipode.** If  $x \in H_\Omega^n(G, k)$  is represented by an  $n$ -fold  $k$ -squeezed extension (4.10), then we can form a dual extension

$$0 \rightarrow k \rightarrow M_0^* \rightarrow \cdots \rightarrow M_{n-1}^* \rightarrow k \rightarrow 0.$$

Here,  $M^* = \text{Hom}_k(M, k)$  is the  $k$ -linear dual of the  $kG$ -module  $M$ . Let  $\tau(x) \in H_\Omega^n(G, k)$  be the class of this extension. Then  $\tau: H_\Omega^n(G, k) \rightarrow H_\Omega^n(G, k)$  is an antiautomorphism of order two for both the product and the coproduct.

The coproduct (4.4), product (4.7) and antipode (4.11) make  $H_\Omega^*(G, k)$  into a commutative (but not cocommutative) Hopf algebra.

### 5. HOMOLOGICAL REFORMULATION

In this section we describe the homology of  $k$ -squeezed resolutions in terms of Tor and Ext over a finite dimensional algebra closely related to the group algebra. Let  $f$  be an idempotent in  $kG$  such that  $f.kG$  is the projective cover of the trivial module, and let  $e = 1 - f$ . Let  $A$  denote the algebra  $e.kG.e$ . The following is Theorem 1.1 from the Introduction.

**Theorem 5.1.** (i) We have isomorphisms  $H_n^\Omega(G, k) \cong \text{Tor}_{n-1}^A(kG.e, e.kG)$  for  $n \geq 2$  and an exact sequence

$$0 \rightarrow H_1^\Omega(G, k) \rightarrow kG.e \otimes_A e.kG \rightarrow kG \rightarrow H_0^\Omega(G, k) \rightarrow 0.$$

(ii) We have isomorphisms  $H_{\Omega}^n(G, k) \cong \text{Ext}_A^{n-1}(e.kG, e.kG)$  for  $n \geq 2$  and an exact sequence

$$0 \rightarrow H_{\Omega}^0(G, k) \rightarrow kG \rightarrow \text{Hom}_A(e.kG, e.kG) \rightarrow H_{\Omega}^1(G, k) \rightarrow 0.$$

*Proof.* (i) Given a projective resolution of  $e.kG$  as an  $A$ -module,

$$\dots \rightarrow \bar{P}_2 \rightarrow \bar{P}_1 \rightarrow \bar{P}_0 \rightarrow 0,$$

we consider the complex of  $kG$ -modules

$$(5.2) \quad \dots \rightarrow kG.e \otimes_A \bar{P}_2 \rightarrow kG.e \otimes_A \bar{P}_1 \rightarrow kG.e \otimes_A \bar{P}_0 \rightarrow kG \rightarrow 0$$

where the last non-zero map is the composite of the augmentation map and the multiplication map

$$kG.e \otimes_A \bar{P}_0 \rightarrow kG.e \otimes_A e.kG \rightarrow kG.$$

It is not hard to show that (5.2) is a left  $k$ -squeezed resolution. Define  $P_0 = kG$  and  $P_i = kG.e \otimes_A \bar{P}_{i-1}$  for  $i \geq 1$ . Then  $P_*$  is the mapping cone of the morphism of complexes  $kG.e \otimes_A \bar{P}_* \rightarrow kG$ . The long exact sequence of this mapping cone gives the required statements.

(ii) Dually, given a weakly injective resolution of  $e.kG$  by projective  $A$ -modules,

$$0 \rightarrow \bar{I}_0 \rightarrow \bar{I}_1 \rightarrow \bar{I}_2 \rightarrow \dots,$$

we consider the complex of  $kG$ -modules

$$(5.3) \quad 0 \rightarrow kG \rightarrow \text{Hom}_A(e.kG, \bar{I}_0) \rightarrow \text{Hom}_A(e.kG, \bar{I}_1) \rightarrow \text{Hom}_A(e.kG, \bar{I}_2) \rightarrow \dots,$$

where the first map is the composite

$$kG \rightarrow \text{Hom}_A(e.kG, e.kG) \rightarrow \text{Hom}_A(e.kG, \bar{I}_0).$$

It is not hard to show that (5.3) is a right  $k$ -squeezed resolution. Define  $I_0 = kG$  and  $I_{-i} = \text{Hom}_A(e.kG, \bar{I}_{-i+1})$  for  $i \geq 1$ . Then  $I_*$  is the mapping cone of the morphism of complexes  $kG \rightarrow \text{Hom}_A(e.kG, \bar{I}_*)$ . The long exact sequence of this mapping cone again gives the required statements.  $\square$

### 6. LOOPS ON $BG_p^\wedge$

In this section, we give a homotopy theoretic interpretation of the squeezed resolutions described in Section 3. We begin by recalling that the Bousfield–Kan completion functor of [5] with respect to the coefficient ring  $\mathbb{F}_p$  takes  $BG$  to a space which we denote  $BG_p^\wedge$ . There is a map  $BG \rightarrow BG_p^\wedge$  which is a mod  $p$  homology equivalence, and we have  $\pi_1(BG_p^\wedge) \cong G/O^p(G)$ . Applying the Hurewicz theorem to the universal cover of  $BG_p^\wedge$ , we see that  $\pi_2(BG_p^\wedge) \cong H_2(O^p(G), \mathbb{Z}_p)$ .

Denote by  $A_p(G)$  the homotopy fibre of the completion map  $BG \rightarrow BG_p^\wedge$ , and choose a basepoint in  $A_p(G)$  mapping to the basepoint in  $BG$ . Then the long exact sequence of the fibration shows that there is an exact sequence

$$1 \rightarrow H_2(O^p(G), \mathbb{Z}_p) \rightarrow \pi_1(A_p(G)) \rightarrow O^p(G) \rightarrow 1,$$

and since  $A_p(G)$  is  $\mathbb{F}_p$ -acyclic, it follows that this is the universal  $p$ -central extension of  $O^p(G)$ .

Now form the pullback  $L_p(G)$  as in the following diagram:

$$\begin{array}{ccccccc} & & L_p(G) & \longrightarrow & EG & & \\ & & \downarrow & & \downarrow & & \\ \Omega BG_p^\wedge & \longrightarrow & A_p(G) & \longrightarrow & BG & \longrightarrow & BG_p^\wedge \end{array}$$

Thus  $L_p(G)$  is homotopy equivalent to  $\Omega BG_p^\wedge$ . The advantage of the space  $L_p(G)$ , though, is that it comes with a free action of  $G$  with connected acyclic quotient  $A_p(G)$ . So the chains  $C_*(L_p(G); k)$  and the cochains  $C^*(L_p(G); k)$  form complexes of free  $kG$ -modules.

**Theorem 6.1.**  $C_*(L_p(G); k)$  is a left  $k$ -squeezed resolution for  $G$ , and  $C^*(L_p(G); k)$  is a right  $k$ -squeezed resolution for  $G$ .

*Proof.* The fact that  $G$  acts freely on  $L_p(G)$  gives condition (i) in Definitions 3.2 and 3.3. The fact that the quotient is connected and acyclic gives condition (ii). The fact that  $O^p(G)$  acts trivially on  $H_*(L_p(G); k)$  and on  $H^*(L_p(G); k)$  gives condition (iii).  $\square$

**Corollary 6.2.** *There are natural isomorphisms*

$$\begin{aligned} H_*^\Omega(G, k) &\cong H_*(\Omega BG_p^\wedge; k), \\ H_\Omega^*(G, k) &\cong H^*(\Omega BG_p^\wedge; k). \end{aligned}$$

*Proof.* This follows from Definition 3.6 and Theorem 6.1.  $\square$

This completes the proof of Theorem 1.2 of the Introduction.

### 7. APPROXIMATION IN $\mathbf{Klnj}(kG)$

In this and the next section, we give another way of viewing the translation between the topology and the algebra.

We write  $\mathbf{Klnj}(kG)$  for the homotopy category of chain complexes of injective (or equivalently, projective)  $kG$ -modules. This category was studied in detail in Benson and Krause [2]. We recall some definitions from Dwyer and Greenlees [7], adapted to this context. Let  $ik$  denote an injective resolution of  $k$ , regarded as an object in  $\mathbf{Klnj}(kG)$ , where it is well defined up to isomorphism. In  $\mathbf{Klnj}(kG)$ , we say that an object  $N$  is *ik-trivial* if the chain complex of homomorphisms  $\mathrm{Hom}_{kG}(ik, N) \simeq 0$ . A map is an *ik-equivalence* if its mapping cone is *ik-trivial*. We say that  $X$  is *ik-torsion* if we have  $\mathrm{Hom}_{kG}(X, N) \simeq 0$  for all *ik-trivial*  $N$ . We say that  $X$  is *ik-complete* if we have  $\mathrm{Hom}_{kG}(N, X) \simeq 0$  for all *ik-trivial*  $N$ . We write  $ik\text{-Tors}(kG)$  and  $ik\text{-Comp}(kG)$  for the subcategories of *ik-torsion* and *ik-complete* complexes.

*Remark 7.1.* It is shown in Theorem 12.3 of [2] that localising subcategories of  $\mathbf{Klnj}(kG)$  are closed under filtered colimits. From this, it can be easily deduced that an object  $X$  in  $\mathbf{Klnj}(kG)$  is *ik-torsion* if and only if it is in the localising subcategory generated by  $ik$ .

Given any object  $X$  in  $\mathbf{Klnj}(kG)$ , there exists an *ik-torsion* object  $\mathrm{Cell}_{ik}(X)$  and an *ik-equivalence*  $\mathrm{Cell}_{ik}(X) \rightarrow X$ . Such an object and map are unique up to isomorphism.

Similarly, given any object  $X$  in  $\mathbf{Klnj}(kG)$ , there exists an *ik-complete* object  $X_{ik}^\wedge$  and an *ik-equivalence*  $X \rightarrow X_{ik}^\wedge$ . This is also unique up to isomorphism.

**Lemma 7.2.** (i) Let  $I_*$  be an object in  $\text{Klnj}(kG)$  which is bounded above and whose homology satisfies  $[OP(G), H_*(I_*)] = 0$ . Then  $I_*$  is  $ik$ -torsion.

(ii) Let  $P_*$  be an object in  $\text{Klnj}(kG)$  which is bounded below and whose homology satisfies  $[OP(G), H_*(P_*)] = 0$ . Then  $P_*$  is  $ik$ -complete.

*Proof.* (i) Let  $n$  be the largest integer such that  $H_n I_* \neq 0$ . Since  $I_*$  is bounded above, it is equivalent to a complex  $I(0)_*$  such that  $I(0)_j = 0$  for  $j > n$ . Let  $X(0)_*$  be an injective resolution of  $\text{Soc} H_n I(0)_*$ , shifted in degree by  $n$ , so that  $X(0)_*$  is a direct sum of copies of shifts of  $ik$ . Then the isomorphism  $H_n X(0)_* \rightarrow \text{Soc} H_n I(0)_*$  extends to a map of complexes  $X(0)_* \rightarrow I(0)_*$ . Write  $I(1)_*$  for the mapping cone, so that we have a triangle  $X(0)_* \rightarrow I(0)_* \rightarrow I(1)_*$ . Now  $I(1)_*$  has smaller homology than  $I(0)$ , in the sense that either  $H_j I(1)_* = 0$  for  $j \geq n$  or  $H_j I(1)_* = 0$  for  $j > n$ , and the radical length of  $H_n I(1)_*$  is smaller than that of  $H_n I(0)_*$ .

So we repeat and get  $X(1)_* \rightarrow I(1)_* \rightarrow I(2)_*$ , and so on. By the octahedral axiom, we have an object  $Y(1)_*$  and triangles  $X(0)_* \rightarrow Y(1)_* \rightarrow X(1)_*$  and  $Y(1)_* \rightarrow I(0)_* \rightarrow I(2)_*$ . Proceeding inductively with  $Y(0)_* = X(0)_*$ , we obtain objects  $X(i)_*$ ,  $Y(i)_*$  and  $I(i)_*$  fitting into commutative diagrams

$$\begin{array}{ccccc}
 Y(i-1)_* & \xlongequal{\quad} & Y(i-1)_* & & \\
 \downarrow & & \downarrow & & \\
 Y(i)_* & \longrightarrow & I(0)_* & \longrightarrow & I(i+1)_* \\
 \downarrow & & \downarrow & & \downarrow \\
 X(i)_* & \longrightarrow & I(i)_* & \longrightarrow & I(i+1)_*
 \end{array}$$

Taking the homotopy colimit of the sequence of triangles

$$\begin{array}{ccccccc}
 Y(0)_* & \longrightarrow & Y(1)_* & \longrightarrow & Y(2)_* & \longrightarrow & \cdots \\
 \downarrow & & \downarrow & & \downarrow & & \\
 I(0)_* & \xlongequal{\quad} & I(0)_* & \xlongequal{\quad} & I(0)_* & \xlongequal{\quad} & \cdots \\
 \downarrow & & \downarrow & & \downarrow & & \\
 I(1)_* & \longrightarrow & I(2)_* & \longrightarrow & I(3)_* & \longrightarrow & \cdots
 \end{array}$$

we have  $\varinjlim_i I(i)_* \simeq 0$ , and so  $I_* \simeq I(0)_* \simeq \varinjlim_i Y(i)_*$  is in the localising subcategory of  $\text{Klnj}(kG)$  generated by  $ik$ .

(ii) The proof of this is dual. There is just one more point to be made, which is that generally the limit of an inverse system of triangles is not a triangle. However, the inverse systems in question are eventually constant in any particular degree, so the inverse limit of the triangles is indeed a triangle.  $\square$

**Theorem 7.3.** (i) If  $I_*$  is a right  $k$ -squeezed resolution, then  $I_* \cong \text{Cell}_{ik}(kG)$  in  $\text{Klnj}(kG)$ .

(ii) If  $P_*$  is a left  $k$ -squeezed resolution, then  $P_* \cong kG_{ik}^\wedge$  in  $\text{Klnj}(kG)$ .

*Proof.* (i) Lemma 7.2 (i) shows that  $I_*$  is  $ik$ -torsion. There is a map  $I_* \rightarrow kG$  which is an  $ik$ -equivalence, because its mapping cone is equivalent to a complex consisting of injective modules which admit no homomorphisms from  $k$ .

(ii) Lemma 7.2 (ii) shows that  $P_*$  is  $ik$ -complete. There is a map  $kG \rightarrow P_*$  which is an  $ik$ -equivalence for the same reason as before.  $\square$

Now recall that if  $X$  is simply connected or  $k$  is a field of characteristic  $p$  and  $\pi_1(X)$  is a finite  $p$ -group, then the Eilenberg–Moore construction gives equivalences

$$\begin{aligned} \mathbb{R}\text{End}_{C^*(X;k)}(k) &\cong C_*(\Omega X; k), \\ k \otimes_{C^*(X;k)}^{\mathbb{L}} k &\cong C^*(\Omega X; k) \end{aligned}$$

(see §4.22 of [8]). Here,  $\mathbb{R}\text{End}$  and  $\otimes^{\mathbb{L}}$  denote the total derived functors of  $\text{End}$  and  $\otimes$ . We can't apply this directly to  $BG$  with  $G$  a finite group, because  $\pi_1(BG) = G$  is not necessarily a finite  $p$ -group. However,  $\pi_1(BG_p^\wedge) = G/O^p(G)$  is a finite  $p$ -group, and we have  $C^*(BG_p^\wedge; k) \simeq C^*(BG; k)$ . In particular, we have

$$(7.4) \quad C_*(\Omega BG_p^\wedge; k) \cong \mathbb{R}\text{End}_{C^*(BG;k)}(k),$$

$$(7.5) \quad C^*(\Omega BG_p^\wedge; k) \cong k \otimes_{C^*(BG;k)}^{\mathbb{L}} k.$$

Let  $\mathcal{E}_G$  be the differential graded algebra  $\text{End}_{kG}(ik)$ . Recall from Theorem 3.3 and Lemma 6.4 of Benson and Krause [2] that there is an adjunction

$$\text{Klnj}(kG) \begin{array}{c} \xleftarrow{-\otimes_{\mathcal{E}_G}^{\mathbb{L}} ik} \\ \xrightarrow{\text{Hom}_{kG}(ik, -)} \end{array} \text{D}_{\text{dg}}(\mathcal{E}_G) \simeq \text{D}_{\text{dg}}(C^*(BG; k))$$

inducing an equivalence between  $\text{D}_{\text{dg}}(C^*(BG; k))$  and  $ik\text{-Tors}(kG)$ . The composite is the “cellularisation functor” in  $\text{Klnj}(kG)$ :

$$\text{Cell}_{ik}(X) = \text{Hom}_{kG}(ik, X) \otimes_{\mathcal{E}_G}^{\mathbb{L}} ik.$$

We have  $\text{Hom}_{kG}(ik, kG) \cong k$  in  $\text{D}_{\text{dg}}(\mathcal{E}_G)$ , so that under this correspondence, the object  $k$  in  $\text{D}_{\text{dg}}(C^*(BG; k))$  corresponds to  $\text{Cell}_{ik}(kG)$  in  $ik\text{-Tors}(kG)$ .

Similarly, the functor  $\text{Hom}_{kG}(ik, -): \text{Klnj}(kG) \rightarrow \text{D}_{\text{dg}}(C^*(BG; k))$  has a right adjoint, and this induces an equivalence between  $\text{D}_{\text{dg}}(C^*(BG; k))$  and  $ik\text{-Comp}(kG)$ . The composite is the “completion functor” in  $\text{Klnj}(kG)$  sending  $X$  to  $X_{ik}^\wedge$ .

**Theorem 7.6.** *We have*

- (i)  $C_*(\Omega BG_p^\wedge; k) \cong \text{End}_{kG}(\text{Cell}_{ik}(kG)) \cong \text{End}_{kG}(kG_{ik}^\wedge)$ ,
- (ii)  $H_*(\Omega BG_p^\wedge; k) \cong H_*(kG_{ik}^\wedge)$ ,
- (iii)  $H^*(\Omega BG_p^\wedge; k) \cong H_{-*}(\text{Cell}_{ik}(kG))$ .

*Proof.* Part (i) follows from the previous discussion.

(ii) Using the map  $kG \rightarrow kG_{ik}^\wedge$ , we have isomorphisms

$$H_*(kG_{ik}^\wedge) \cong \text{Hom}_{kG}(kG, kG_{ik}^\wedge)_* \cong \text{Hom}_{kG}(kG_{ik}^\wedge, kG_{ik}^\wedge)_*.$$

(iii) follows from (ii) by dualising both sides.  $\square$

Combining Theorems 7.3 and 7.6, we obtain our second proof of Theorem 1.2. The advantage of this proof is that it makes it clear that the product (4.7) in  $H_*^\Omega(G, k)$  corresponds to the loop product in  $H_*(\Omega BG_p^\wedge; k)$ . This does not seem so apparent from the proof in Section 6.

## 8. FINAL REMARKS

Using work of Félix, Halperin and Thomas [12], Levi [19] proves that  $H_*(\Omega BG_p^\wedge; k)$  has either polynomial or semiexponential growth. Here, semiexponential growth means that there exists  $\lambda > 1$  such that for an infinite number of positive integers  $n$  the dimension of the  $n$ th homology group is at least  $\lambda^{\sqrt{n}}$ . In the case of polynomial growth, the homology is finitely generated, nilpotent, and a finite module over a central polynomial subalgebra (cf. Theorem A of [10], Theorem B of [11]). No examples are known where the growth is semiexponential but not exponential. We close with some remarks on the boundary between polynomial and semiexponential growth.

A theorem of Gulliksen [14, 15] states that if  $R$  is a commutative Noetherian local ring with residue field  $k$ , then  $R$  is a complete intersection if and only if  $\text{Ext}_R^*(k, k)$  has polynomial growth.

In the case of connected graded commutative algebras, we have to modify the definition slightly to take into account the fact that odd degree elements anticommute instead of commute. We say that a graded commutative ring  $R$  is of *polynomial type* over  $R_0 = k$  if it can be generated by a set of elements  $x_1, \dots, x_r$  of positive degree subject only to the relations coming from graded commutativity. Notice that if  $x_i$  is an element of odd degree, then graded commutativity forces  $2x_i^2 = 0$ . If  $k$  is a field of characteristic two, then this definition just gives polynomial rings. If  $k$  is a field of odd characteristic, on the other hand, then it defines a polynomial ring on the even degree generators tensored with an exterior algebra on the odd degree generators.

A homogeneous element  $y$  of a graded commutative ring  $R$  is said to be *regular* if the annihilator of  $y$  is  $(y - (-1)^{|y|}y)$ ; i.e., it is the zero ideal if  $|y|$  is even and  $(2y)$  if  $|y|$  is odd. A sequence of elements  $y_1, \dots, y_s$  of positive degree is said to be a *regular sequence* if each  $y_i$  is regular in  $R/(y_1, \dots, y_{i-1})$ .

A connected graded  $k$ -algebra is said to be a *complete intersection* if it can be written as a quotient of a  $k$ -algebra of polynomial type by a regular sequence. With this definition, Gulliksen's theorem continues to hold (Avramov and Iyengar, unpublished).

**Theorem 8.1.** *If  $H^*(G, k)$  is a complete intersection, then  $H_*(\Omega BG_p^\wedge; k)$  has polynomial growth.*

*Proof.* The theorem follows from Gulliksen's theorem and the Eilenberg–Moore spectral sequence

$$\text{Ext}_{H^*(G, k)}^{**}(k, k) \Rightarrow H_*(\Omega BG_p^\wedge; k). \quad \square$$

We can prove the converse of Theorem 8.1 in the special case of an elementary abelian Sylow 2-subgroup in characteristic two. In this case, there is no ambiguity as to what is meant by a complete intersection since there are no signs to worry about.

**Theorem 8.2.** *Suppose that  $k$  is a field of characteristic two, and that  $G$  has an elementary abelian Sylow 2-subgroups  $E$ . Then the following are equivalent:*

- (i)  $H_*(\Omega BG_2^\wedge; k)$  has polynomial growth.
- (ii)  $H^*(G, k)$  is a complete intersection.
- (iii)  $N_G(E)/C_G(E)$  is an odd order group generated by elements of order three whose fixed point set on  $E$  has index four.

(iv)  $N_G(E)/O_{2'}N_G(E)$  is a direct product of copies of  $\mathbb{Z}/2$ ,  $A_4 = (\mathbb{Z}/2)^2 \rtimes \mathbb{Z}/3$  and  $(\mathbb{Z}/2)^3 \rtimes F_{21}$ , where  $F_{21}$  is the non-abelian group of order 21 and the action on  $(\mathbb{Z}/2)^3$  is faithful.

(v)  $G/O_{2'}(G)$  is a direct product of copies of  $\mathbb{Z}/2$ ,  $A_4$ ,  $L_2(q)$  ( $q = 4$  or  $q \equiv 3, 5 \pmod{8}$ ), the Ree groups  ${}^2G_2(3^{2m+1})$  ( $m \geq 1$ ),  $(\mathbb{Z}/2)^3 \rtimes F_{21}$ ,  $L_2(8) \rtimes \mathbb{Z}/3$  and the Janko group  $J_1$ .

*Proof.* First we prove that (i) is equivalent to (ii). If  $G$  has an elementary abelian Sylow 2-subgroup, then  $C^*(BG; k)$  is equivalent as a differential graded algebra to its cohomology. To see this, first it's true for  $G = \mathbb{Z}/2$  since the reduced bar complex is the minimal resolution. For a more general elementary abelian 2-group, we use tensor products of copies of the reduced bar complex for the direct factors of the group to see that  $C^*(BG; k)$  is equivalent to  $H^*(G, k)$ . In the case of a group with elementary abelian Sylow 2-subgroup  $E$ ,  $C^*(BG; k)$  is equivalent to the invariants of  $N_G(E)/C_G(E)$  on  $H^*(E, k)$ .

It follows that the Eilenberg–Moore spectral sequence stops at the  $E_2$  page, so that the dimension of  $H_n(\Omega BG_p^\wedge; k)$  is equal to  $\sum_{i+j=n} \dim_k \text{Ext}_{H^*(G, k)}^{i, j}(k, k)$ . Now we can use Gulliksen's theorem to see that (i) is equivalent to (ii).

Since  $H^*(G, k) = H^*(E, k)^{N_G(E)/C_G(E)}$ , the question of when (ii) holds reduces to the question of when the polynomial invariants of an odd order group over  $\mathbb{F}_2$  form a complete intersection.

To prove that (ii) implies (iii), we remark that the general question of when polynomial invariants of a finite group form a complete intersection was studied by Gordeev [13], Kac and Watanabe [16] and Nakajima [21, 22, 23], among others. In particular, it is known that if the invariants form a complete intersection, then the group is generated by elements whose fixed space has codimension at most two. Translating this into group theory, this means that  $N_G(E)/C_G(E)$  should be generated by elements of order three whose fixed points on  $E$  have index four. This completes the proof that (ii) implies (iii).

The statement that (iv) implies (ii) is just a question of checking that the cohomology is a complete intersection in the cases given and then using the Künneth formula for the cohomology of a product. The equivalence of (iv) and (v) is easy to deduce from John Walter's classification [24] of finite groups with an abelian Sylow 2-subgroup, together with Bombieri's proof [3] that the groups of Ree type are Ree groups.

It remains to prove that (iii) implies (iv). I am grateful to Geoff Robinson for supplying me with a proof of this. We use the phrase "3-reflection" for an element of order three in  $\bar{G} = N_G(E)/C_G(E)$  whose centraliser in  $E$  has index four.

First we show that we may decompose  $\bar{G}$  as a direct product of groups  $\bar{G}_i$ , and  $E$  as a direct product of  $E_i$ , so that  $\bar{G}_i$  acts trivially on  $E_j$  for  $j \neq i$ , and  $\bar{G}_i$  is generated by a single conjugacy class of 3-reflections. To see this, we note that if  $x$  and  $y$  are 3-reflections in  $\bar{G}$  which generate non-conjugate subgroups, then  $\langle x, y \rangle$  is isomorphic to an odd order subgroup of  $\text{GL}(4, 2)$  which is not isomorphic to  $F_{21}$  and which must therefore be an elementary abelian 3-group. It follows that the conjugates of  $x$  and the conjugates of  $y$  generate mutually centralising normal subgroups of  $\bar{G}$ . Suppose that  $U$  is an irreducible summand of  $E$  on which both  $x$  and  $y$  act non-trivially. Setting  $\mathbb{F} = \text{End}_{\mathbb{F}_2 \bar{G}}(U)$ , we can regard  $U$  as an  $\mathbb{F}\bar{G}$ -module. We can write  $U = U_1 \otimes_{\mathbb{F}} U_2$  so that  $x$  acts as scalars on  $U_1$  and non-trivially on  $U_2$ , and  $y$  acts as scalars on  $U_2$  and non-trivially on  $U_1$ . Since  $x$  and  $y$  are 3-reflections,

this implies that either  $U_1$  or  $U_2$  is one dimensional; then we deduce that the other is also one dimensional. So  $x$  and  $y$  both act as scalars and  $U$  is one dimensional over  $\mathbb{F}$ . Thus  $y$  acts as  $x$  or  $x^{-1}$  on  $U$  and trivially on the remaining summands of  $E$ . So we can remove  $y$  from the generating set without loss. This completes the proof of the decomposition.

This means that we may assume that the action of  $\bar{G}$  on  $E$  is faithful and irreducible. If  $\bar{G}$  is trivial, then  $E \cong \mathbb{Z}/2$ . Assuming we are not in this case,  $\bar{G}$  is generated by a single conjugacy class of 3-reflections. The next step is to show that  $E$  is primitive. If it is induced from an  $\mathbb{F}_2 H$ -module  $V$  for a proper subgroup  $H$  of  $\bar{G}$ , then  $x$  does not lie in the intersection of the conjugates of  $H$ , so  $V$  must be one dimensional, since otherwise  $x$  could not act as a 3-reflection. But then  $V$  is the trivial module, and this contradicts the irreducibility of  $E$ . Thus  $E$  is primitive.

As before, let  $\mathbb{F} = \mathbb{F}_{2^e} = \text{End}_{\mathbb{F}_2 \bar{G}}(E)$ , so that we may regard  $E$  as an  $\mathbb{F} \bar{G}$ -module. Since  $\bar{G}$  contains a 3-reflection  $x$ , we have  $e = 1$  or  $2$ . If  $e = 2$ , then the fixed points of  $x$  on  $E$  have  $\mathbb{F}$ -codimension one. So for any  $g \in \bar{G}$ ,  $\langle x, x^g \rangle$  is isomorphic to an odd order subgroup of  $GL(2, 4) \cong \mathbb{Z}/3 \times A_5$  generated by elements of order three. Hence  $\langle x, x^g \rangle$  is abelian. Since  $\bar{G}$  is generated by the conjugates of  $x$ , it follows that  $\bar{G}$  is cyclic, so  $N_G(E)/O_2 N_G(E) \cong A_4$  and  $E \cong (\mathbb{Z}/2)^2$ . So if we are not in this case, we may assume that  $E$  is absolutely irreducible.

If  $E$  were absolutely primitive as an  $\mathbb{F}_2 \bar{G}$ -module, then using Clifford theory with respect to the Fitting subgroup  $F(\bar{G})$  of the soluble group  $\bar{G}$ , we see that  $C_{\bar{G}}(F(\bar{G})) = ZF(\bar{G})$  would be central in  $\bar{G}$ , and hence by Schur's lemma it would be trivial, a contradiction. So for some finite extension  $\mathbb{F}$  of  $\mathbb{F}_2$ ,  $\mathbb{F} \otimes_{\mathbb{F}_2} E$  is induced from some proper maximal subgroup  $H$  of  $\bar{G}$ . If  $x$  is a 3-reflection in  $\bar{G}$ , then  $x$  must act as a single 3-cycle and possibly some fixed points on the cosets of  $H$ , or else the eigenvalues would be wrong. So the permutation action of  $\bar{G}$  on the cosets of  $H$  is generated by 3-cycles and has odd order, so these 3-cycles commute. So  $G$  modulo the intersection of the conjugates of  $H$  is abelian, and hence  $H$  is normal of index 3. Since  $\bar{G}$  contains 3-reflections, it follows that it's induced from a one dimensional representation of  $H$ , and so  $E$  is three dimensional. Thus  $\bar{G}$  is isomorphic to a subgroup of  $GL(3, 2)$ , and the only possibility is  $\bar{G} \cong F_{21}$ .  $\square$

*Remark 8.3.* The smallest simple group with elementary abelian Sylow 2-subgroups, and not satisfying the conditions of the theorem, is  $L_2(8)$ . Since the Sylow 2-normalizer in this group is  $(\mathbb{Z}/2)^3 \rtimes \mathbb{Z}/7$ , this corresponds to the first row of the table in Section 2.

**Conjecture 8.4.** *If  $G$  has abelian Sylow  $p$ -subgroups, then  $H^*(G, k)$  is a complete intersection if and only if  $H_*(\Omega B G_p^\wedge; k)$  has polynomial growth.*

The conjecture fails badly when the Sylow  $p$ -subgroups are not abelian. For example, there are plenty of  $p$ -groups whose cohomology rings are not complete intersections. In this situation, we should use a "derived" notion of complete intersection. Dwyer, Greenlees and Iyengar [9] formulate the following definition, which we shall call a "quasi-complete intersection".

**Definition 8.5.** A differential graded augmented algebra  $\Lambda \rightarrow k$  is said to be a quasi-complete intersection if, given any object  $X$  in  $D(\Lambda)$  with the augmentation ideal in its support, the thick subcategory it generates contains a non-zero object in the thick subcategory generated by  $\Lambda$ .

The following conjecture is due to John Greenlees:

**Conjecture 8.6.** *For any finite group  $G$ ,  $C^*(BG; k)$  is a quasi-complete intersection if and only if  $H_*(\Omega BG_p^\wedge; k)$  has polynomial growth.*

Finally, we reiterate and strengthen a conjecture made by Levi [17].

**Conjecture 8.7.** *For any finite group  $G$ ,  $H_*(\Omega BG_p^\wedge; k)$  is a finitely generated  $k$ -algebra. Furthermore, either it has polynomial growth or it contains a free algebra on two generators.*

#### ACKNOWLEDGMENT

The author thanks the following people for stimulating conversations about this work: Bill Dwyer, John Greenlees, Henning Krause, Ran Levi, Markus Linckelmann, Geoff Robinson and Raphaël Rouquier.

#### REFERENCES

1. D. J. Benson, *Representations and Cohomology II: Cohomology of groups and modules*, Cambridge Studies in Advanced Mathematics, vol. 31, Cambridge University Press, 1991, reprinted in paperback, 1998. MR1156302 (93g:20099)
2. D. J. Benson and H. Krause, *Complexes of injective  $kG$ -modules*, *Algebra and Number Theory* **2** (2008), 1–30. MR2377361
3. E. Bombieri, *Thompson's problem ( $\sigma^2 = 3$ )*, *Invent. Math.* **58** (1980), 77–100. MR570875 (81f:20019)
4. W. Bosma and J. Cannon, *Handbook of Magma Functions*, Magma Computer Algebra, Sydney, 1996.
5. A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, *Lecture Notes in Mathematics*, vol. 304, Springer-Verlag, Berlin/New York, 1972. MR0365573 (51:1825)
6. F. R. Cohen and R. Levi, *On the homotopy theory of  $p$ -completed classifying spaces*, *Group representations: cohomology, group actions and topology* (Seattle, WA, 1996), *Proc. Symp. Pure Math.*, vol. 63, American Math. Society, 1998, pp. 157–182. MR1603147 (99e:55026)
7. W. G. Dwyer and J. P. C. Greenlees, *Complete modules and torsion modules*, *Amer. J. Math.* **124** (2002), 199–220. MR1879003 (2003g:16010)
8. W. G. Dwyer, J. P. C. Greenlees, and S. Iyengar, *Duality in algebra and topology*, *Adv. in Math.* **200** (2006), 357–402. MR2200850 (2006k:55017)
9. ———, *Finiteness in derived categories of local rings*, *Comment. Math. Helvetici* **81** (2006), 383–432. MR2225632 (2007b:13021)
10. Y. Félix, S. Halperin, and J.-C. Thomas, *Hopf algebras of polynomial growth*, *J. Algebra* **125** (1989), 408–417. MR1018954 (90j:16021)
11. ———, *Elliptic Hopf algebras*, *J. London Math. Soc.* **43** (1991), 545–555. MR1113392 (92i:57033)
12. ———, *Elliptic spaces II*, *Enseign. Math.* **39** (1993), 25–32. MR1225255 (94f:55008)
13. N. L. Gordeev, *Finite linear groups whose algebra of invariants is a complete intersection*, *Izv. Akad. Nauk SSSR Ser. Mat.* **50** (1986), 343–392. MR842586 (88a:13020)
14. T. H. Gulliksen, *A homological characterization of local complete intersections*, *Compositio Math.* **23** (1971), 251–255. MR0301008 (46:168)
15. ———, *On the deviations of a local ring*, *Math. Scand.* **47** (1980), 5–20. MR600076 (82c:13022)
16. V. Kac and K. Watanabe, *Finite linear groups whose ring of invariants is a complete intersection*, *Bull. Amer. Math. Soc.* **6** (1982), 221–223. MR640951 (83h:14042)
17. R. Levi, *On finite groups and homotopy theory*, *Mem. AMS*, vol. 567, American Math. Society, 1995. MR1308466 (96c:55019)
18. ———, *A counter-example to a conjecture of Cohen*, *Algebraic topology: new trends in localization and periodicity* (Sant Feliu de Guixols, 1994), *Birkhäuser Verlag*, Basel, 1996, pp. 261–269. MR1397737 (97d:55027)

19. ———, *On homological rate of growth and the homotopy type of  $\Omega BG_p^\wedge$* , *Math. Zeit.* **226** (1997), 429–444. MR1483541 (98k:55016)
20. ———, *On  $p$ -completed classifying spaces of discrete groups and finite complexes*, *J. London Math. Soc.* **59** (1999), 1064–1080. MR1709097 (2000m:55019)
21. H. Nakajima, *Quotient singularities which are complete intersections*, *Manuscripta Math.* **48** (1984), 163–187. MR753729 (86h:14039)
22. ———, *Quotient complete intersections of affine spaces by finite linear groups*, *Nagoya Math. J.* **98** (1985), 1–36. MR792768 (87c:14056)
23. H. Nakajima and K. Watanabe, *The classification of quotient singularities which are complete intersections*, *Complete Intersections, Acireale 1983*, *Lecture Notes in Mathematics*, vol. 1092, Springer-Verlag, Berlin/New York, 1984, pp. 102–120. MR775879 (86c:14002)
24. J. H. Walter, *The characterization of finite groups with abelian Sylow 2-subgroups*, *Ann. of Math.* **89** (1969), 405–514. MR0249504 (40:2749)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE, SCOTLAND  
E-mail address: `bensondj@maths.abdn.ac.uk`