



25 by storm tides, killed *en masse* by exposing fish to hypoxic conditions in a similar way to  
26 modern water bodies effected by storm tides generated during hurricanes.

27

## 28 **1.0 INTRODUCTION**

29 The fish bed section of the Middle Devonian-aged Achanarras Limestone Member, located in the  
30 Orcadian Basin in NE Scotland, preserves an abundant fish fauna many of which display  
31 excellent preservation. Trewin (1986) provided a detailed description of the fish genera present  
32 and their stratigraphic and faunal distribution. In summary; a wide range of life-habitats is  
33 evidenced by the genera encountered that range from *Dipterus* – an early dipnoan (lungfish)  
34 tolerant of dry-intervals that prevailed in shallow water, through to *Coccosteus* the fossils of  
35 which are most abundant in intervals deposited when lakewater levels were high, and which in  
36 many other settings are found in marine environments. Fossil fish are not evenly distributed  
37 throughout the Achanarras Limestone Member but instead found clustered within distinct mass  
38 mortality horizons, with a frequency exceeding one per ten year interval. Hamilton and Trewin  
39 (1988) suggested that mass mortality horizons within the Achanarras Limestone Member could  
40 have resulted from the overturning of a stratified lake and the consequent reduction of oxygen in  
41 surface water as it mixed with hypoxic bottom water. Decay of organic matter generated by algal  
42 blooms stimulated by the resupply of nutrients to surface waters could have further  
43 deoxygenated waters. Although such kill-mechanisms are documented from East African Rift  
44 Lakes (Beadle 1981), no further evidence has been presented to support this mechanism or  
45 indeed the responsible trigger.

46

47 The Achanarras lake itself was deposited in a basin bounded by NE-SW trending faults, with the  
48 nearest coastline (based upon the occurrence of demonstrably marine-successions) located to the  
49 South East (Fig 1). More locally, at least to the SW, SE and NW the Caithness flagstones pass  
50 into fluvial, alluvial and Aeolian facies, thus from a first consideration the lake would appear to  
51 have been truly “landlocked”. However, varying levels of land-sea interaction have been  
52 suggested for the beds. Microfaunal evidence for marine incursions into the Orcadian Basin (e.g.  
53 Marshall et al. 1996) has been reported for the Upper Middle-Devonian (Givetian) of Orkney,  
54 e.g. to the north of the of Orcadian lake. Additionally, certain members of the lakes fish fauna  
55 (e.g. *Cocosteus*) are also found within time-equivalent marine successions deposited in the  
56 ancient Rheic Ocean to the south of the Old Red Continent. However, these fish may have  
57 migrated to the lake and their presence in the fishbeds is therefore not conclusive evidence of an  
58 incursion of seawater (Trewin 1986). Thus there exists evidence of a connection to the marine  
59 environment, but it appears inconsistent, and to have only marginally influenced the lakes  
60 environment.

61  
62 In addition to its fish fauna, the Achanarras Limestone Member is also noted for its seasonally  
63 laminated sediments. The Mid-Devonian 3.6 meter thick Achanarras Limestone Member  
64 comprises lamina couplets interpreted by previous authors as non-glacial varves reflecting intra-  
65 annual variation in sedimentation, where two adjacent laminae account for a years sedimentation  
66 (Rayner, 1963; Trewin, 1986). The varve couplets consist of carbonate / clastic pairs; carbonate  
67 laminae comprise a ferroan microdolomitic carbonate phase and clastic laminae are mainly  
68 siliciclastic (Andrews et al. 2010; Othman Wilson, 2012). A number of orders and scales of  
69 cyclicity are recognised within the Achanarras Limestone Member, but most relevant to this

70 study are the very short order cycles corresponding to Schwabe cycles (~11 years) and shorter  
71 order cycles (3 - 8 years) that most likely represent the oscillation of shorter order climatic cells  
72 (Andrews et al., 2010). This accords with the view of many other workers that the cyclic patterns  
73 of sedimentation observed within the Caithness Flagstone group and particularly the Achanarras  
74 Limestone Member strongly reflect climatic forces rather than tectonic processes and rates of  
75 basin infill (Astin et al., 1990; Marshall et al. 1996; Trewin and Thirlwell 2002).

76

77 The purpose of this study is to look for organic geochemical anomalies that will help ascertain  
78 what was unusual about the Orcadian Lake's water column during the deposition of the fish mass  
79 mortality horizons and better understand its depositional environment. By making geochemical  
80 measurements on a lamina by lamina scale a high resolution chronology can be constructed to  
81 explore how environmental conditions changed and led to the deposition of a mass mortality  
82 horizon.

83

## 84 **2.0 METHODS**

85 Samples were collected from the Achanarras Quarry, Caithness (Fig.1). The specimen used for  
86 this study is from the uppermost fish-bearing section of the Achanarras sequence (facies 6 from  
87 Trewin 1986). Elemental (carbon and sulphur), stable isotope data ( $\delta^{13}\text{C}$  for organic and  
88 inorganic carbon) and biomarker data were obtained for twenty four consecutive laminae.  
89 Limited data is also presented for lamina collected from the middle part of the Achanarras  
90 Limestone Member for comparative purposes.

91

92 Samples for geochemical analysis were obtained by micro-drilling discrete lamina from specially  
93 prepared 5 mm thick slabs using a MicroMill system (New Wave Research Ltd); slabs have flat  
94 polished surfaces, perpendicular to lamina and were lightly etched with hydrochloric acid. The  
95 semi-automated MicroMill, which can articulate submicron distances in three degrees of freedom  
96 (x,y,z), was used to sample laminae less than 1 mm thickness. For thick lamina, a drill mounted  
97 on a manually operated stand was used with a binocular microscope facilitating accurate  
98 positioning. For all methods no lubricating fluid was used and drills were operated at medium-  
99 speed to avoid generating excessive friction or long drilling times. Drill powder was removed  
100 from samples by gravity, air and mechanically by a fine brush. All powder was removed from a  
101 working surface prior to “drilling-out” the next laminae. Laminae were removed alternately, with  
102 in-fill drilling used to sample lamina remaining after the first pass of a sample. While working on  
103 a given slab etching and polishing were repeated to provide clean and clear working surfaces and  
104 prevent cross contamination between samples.

105  
106 Extracts were obtained using a modified version of the mini-extraction methods presented in  
107 Bowden et al., (2008); a three stage extraction process was applied to ~100 mg of powder (using  
108 dichloromethane and methanol), after which extracts were combined and then concentrated by  
109 evaporating the solvent under an inert nitrogen atmosphere. Gas Chromatography Mass  
110 Spectrometry (GC-MS) analysis of the extract was performed using an Agilent 6890N GC fitted  
111 with a J&W DB-5 phase 50 m length column (0.25 mm id, 0.25  $\mu$ m film thickness) connected to  
112 a 5975 MSD and a quadruple mass spectrometer operating in SIM mode (dwell time 0.1 s/ion  
113 and ionisation energy 70 eV). Fifteen ions were monitored;  $m/z$  191, 205 and 412 to help  
114 interpret pentacyclic terpanes such as hopanes,  $m/z$  113, 183 and 125 for isoprenoidal

115 hydrocarbons including  $\beta,\beta$ -carotane and  $m/z$  217, 218, 231 and 259 for four ring terapoids such  
116 as steranes, diasteranes and methylsteranes. Samples were injected manually using a  
117 split/splitless injector operating in splitless mode. Temperature program for the GC oven was 80  
118 – 295 °C, holding at 80 °C for two minutes, rising at 10 °C min<sup>-1</sup> for 8 min and then 3 °C min<sup>-1</sup>  
119 and finally holding the maximum temperature for 10 min. Compounds were identified by  
120 comparing retention times to well-characterised materials that served as reference samples. All  
121 concentrations are reported relative to the internal standard. Illustrative chromatograms for the  
122 biomarkers used in the study are shown in Fig. 2 and Fig. 3.

123  
124 Here we focus on  $\beta,\beta$ -carotane, 24-*n*-propylcholestane (C<sub>30</sub> sterane) and gammacerane  
125 hydrocarbon biomarkers because these compounds demonstrate the clearest links to changes in  
126 palaeoenvironment during the interval concerned. From a technical perspective these compounds  
127 were easy to isolate and measure, because of their relative abundance. Instead of making use of  
128  $\beta,\beta$ -carotane as ratio denominated by another biomarker, we report it as micrograms of  $\beta,\beta$ -  
129 carotane per g of sediment. Doing this permits biomarker concentration to be reported per  
130 laminae – e.g. per unit of time, which thus expresses biomarker data as a net burial rate.  
131 Concentrations of  $\beta,\beta$ -carotane are reported relative to an internal standard of D4-cholestane.  
132 The gammacerane index was calculated after the method presented in Peters et al. (2007), with  
133 peak assignments verified by use of the 412 and 205  $m/z$  ion chromatograms. Similarly sterane  
134 parameters were verified by calculating parameters using the 217 and 218  $m/z$  ion  
135 chromatograms. A comparison of the duplicate parameters obtained is presented in  
136 supplementary information 1. Errors for  $\beta,\beta$ -carotane measurement based on duplicate analysis  
137 of extracts are +/- 5.1 %.

138

139 For  $\delta^{13}\text{C}$  carbonate (inorganic carbon) stable isotope analysis, 1-2 mg samples were dissolved  
140 overnight in phosphoric acid at 70 °C. The carbon dioxide that evolved was purged under  
141 positive pressure, and using helium as the transfer gas analysed on an AP2003 mass  
142 spectrometer. Repeat analyses of NBS-18 and internal calcite standards were generally better  
143 than  $\pm 0.2\%$ .

144

145 For  $\delta^{13}\text{C}$  organic determination powders were acid digested (by sequentially exposing samples  
146 overnight to 10 % and then 25 % hydrochloric acid) to remove all inorganic carbon (carbonate).  
147 Samples rinsed with distilled water, dried and weighed into tin capsules. Samples were then  
148 analysed by continuous flow isotope ratio mass spectrometry (CF-IRMS) using a Thermo  
149 Finnigan Delta Plus XP Mass Spectrometer, coupled to a Costech Elemental Analyser (model  
150 ECS 4010). A minimum of 20 mg (equivalent to approximately 0.1 mg carbon) of sample, per  
151 lamination was combusted in a tin capsule for simultaneous determination of carbon isotope  
152 ratios. Three laboratory standards (prepared from gelatine and alanine standard solutions) were  
153 analysed for every 10 samples, allowing instrument drift to be corrected over the course of a 14  
154 hour analytical sequence. Error on replicates is better than 0.2 ‰. Four aliquots (per run) of  
155 Tryptophan, an amino acid, were also analyzed simultaneously in order to calculate the carbon  
156 content of the samples. All stable isotope ratios are expressed in  $\delta$  notation as parts per thousand  
157 (‰) relative to V-PDB and V-SMOW international standards.

158

### 159 **3.0 RESULTS**

160 The following sections describe data initially from the perspective of establishing an  
161 environmental baseline (dashed line representing an average value in Fig. 4), and then from the  
162 perspective of anomalous values associated with the mass mortality horizon (values exceeding a  
163 standard deviation).

164

### 165 *3.1 Environmental Baseline*

166 Total organic carbon is generally quite low with a biannual average of 0.23% (n= 24,  $\sigma$  =  
167 0.06%). This is consistent with a lake environment in which sedimentation rates were known to  
168 be high (diluting organic carbon), and experienced occasional influxes of relatively coarse  
169 grained sediment that were likely fluvial in origin (Trewin 1986). Such inputs of sediment would  
170 have diluted organic carbon content and lowered TOC values – even if net rates of carbon burial  
171 were high. Conversely carbon/sulphur ratios appear low (there is a lot of sulphur with respect to  
172 carbon), especially for a lacustrine environment and indicate that a relatively high amount of  
173 sulphur was fixed within the ancient sediments as sulphide (Leventhal 1979; Berner and  
174 Raiswell 1986); far higher than would be expected for a freshwater lake or even a marine  
175 environment (e.g. they are less than 2.8). The  $\delta^{13}\text{C}_{\text{carb}}$  data average -1.01‰ (n = 23,  $\sigma$  = 0.31‰)  
176 and the  $\delta^{13}\text{C}_{\text{org}}$  values fall within the standard range reported for algal organic matter (-26 to -  
177 42‰, Leng and Marshall 2004). This might be expected, as a significant input of higher plant  
178 organic carbon would be unlikely for a Middle Devonian-aged sediment.

179

180  $\beta,\beta$ -Carotane is notably prominent in all samples (see Fig. 2) and this is a feature observed in the  
181 solvent extractable organic matter obtained from numerous localities around the Orcadian Basin  
182 (Duncan and Hamilton 1988), where it is often the most easily resolved and abundant



183 hydrocarbon-biomarker on gas chromatograms.  $\beta,\beta$ -Carotane (Fig.2) is derived from  $\beta,\beta$ -  
184 carotene by transformation of the unsaturated hydrocarbon precursor during early diagenesis (cf.  
185 Killops and Killops, 2005). Although  $\beta,\beta$ -carotene is ubiquitous, high concentrations of  $\beta,\beta$ -  
186 carotane in the geological record are not common and the very large proportions of  $\beta,\beta$ -carotane  
187 present in the Achanarras Limestone Member are notable because carotenoids typically degrade  
188 rapidly in most aquatic depositional settings (Jiang and Fowler 1985). Therefore, the very high  
189 proportions of this compound present in the Achanarras Limestone Member and similar  
190 lacustrine rocks and sediments have been interpreted as a consequence of a higher than typical  
191 input from precursor biological materials and a high net primary productivity (Killops and  
192 Killops 2005). Likely sources for this carotenoid-enriched organic matter include halophilic  
193 archaeobacteria which thrive in hypersaline environments (Kushwaha et al. 1974; Rønnekleiv  
194 and Liaaen-Jensen 1996) and contain very high proportions of carotenoids, including  $\beta,\beta$ -  
195 carotene. The concentration of  $\beta,\beta$ -carotane is high in all samples although there are several  
196 instances of values less than one standard deviation from the mean value.

197  
198 The most distinctive feature of the sterane biomarkers is the low abundance of regular  $C_{27}$   
199 steranes (Fig. 3). This likely indicates a low proportion of cholesterol in precursor organic  
200 matter, and hence limited contributions from animals/zoo plankton, which are the main sources  
201 of cholesterol in modern lake sediments (Huang and Meinschein 1979; Kodner et al., 2008). The  
202 24-*n*-propylcholestane ( $C_{30}$  sterane) sterane-homologue is less commonly reported in solvent  
203 extracts obtained from Orcadian Basin sedimentary rocks (Duncan and Hamilton 1988) although  
204 it can be seen to be present in all of the samples during this study, but in varying proportions  
205 (Fig. 3). Regular  $C_{30}$  steranes (24-*n*-propylcholestanes), likely derive from  $C_{30}$  24-*n*-

206 propylcholesterols which have been found to be present in a few largely marine chrysophyte  
207 algae (Rohmer et al., 1980; Moldowan 1984; Volkman 2002). Most important of these is  
208 probably the brown tide alga *Aureococcus anophagefferens* (Giner et al., 2003). Baseline values  
209 for the relative proportion of C<sub>30</sub> sterane for the studied interval are low, both in comparison to  
210 stratigraphically lower intervals of the Achanarras Limestone Member (values shown as crosses  
211 on graph) where the most diverse fish fauna are preserved and also relative to the mass mortality  
212 horizon itself.

213

214 Gammacerane is a pentacyclic triterpanoid hydrocarbon that can be measured on the *m/z* 191  
215 chromatogram (Fig.2) and is present in all samples. The varied proportion found in samples is  
216 indicated by the Gammacerane Index (GI) which is plotted in Fig. 4h. Gammacerane can be used  
217 as an indicator for water column stratification (Sinninghe Damsté et al. 1995; Stephens and  
218 Carroll, 1999). This is because its main biological precursor is tetrahymanol (Ten Haven et al.  
219 1989) a compound that is synthesised by bacterivorous ciliates (Sinninghe Damsté et al. 1995)  
220 inhabiting anoxic waters. (Tetrahymanol is only produced by these organisms in the absence of  
221 dietary sterols, a situation that occurs in the anoxic part of a stratified water column where the  
222 growth of sterol-synthesising eukaryotic algae is inhibited). No precise definition has been  
223 offered as to what constitutes a 'high' GI value, however quoted GI values greater than 0.1 – 0.2  
224 are generally described as being 'high' (e.g. Chen et al. 1996), thus background values indicate  
225 prevailing water column stratification.

226

227 *3.2 Anomalous values associated with the mass mortality horizon*

228 A sharp spike in the C/S ratio two standard deviations high, corresponding to the highest value in  
229 a 10 year interval, occurs during the mass mortality horizon (Fig. 4a – C/S ratio). This is  
230 accompanied by a TOC spike (Fig. 4b) two standard deviations above the average (also a 10 year  
231 maximum) and a reduced burial of sulphur relative to carbon (Fig. 4a), probably indicating less  
232 saline waters or a water column less able to support pyrite formation via reduction of sulphate.  
233 Immediately following the mass mortality horizon there is a large excursion of the  $\delta^{13}\text{C}_{\text{org}}$   
234 parameter (Fig. 4d) to its highest value in an 11 year period of  $-28.65\%$  (average =  $-30.74\%$ ,  $n$   
235 = 24,  $\sigma = 0.93\%$ ). This is still consistent with algal organic matter being the dominant  
236 contributor to the lakes productivity. The concentration of  $\beta,\beta$ -carotane (Fig. 4e) is at a 4 year  
237 low during the mass mortality horizon and then immediately rises to a five year high in the  
238 following year. However, this is one of four big switches (where a parameter changes from a  
239 maximum to minimum value) in this parameter over the 12 year period concerned, and is only  
240 significant because it coincidences with the mass mortality horizon, rather than for its absolute  
241 magnitude (Fig. 4e). The beginning of the mass mortality horizon itself is characterised by a high  
242  $\text{C}_{30}/\text{C}_{28}$  sterane ratio (nearly two standard deviations high) indicating an enhanced burial of  
243 biomarkers derived from marine phytoplankton (Fig. f). The gammacerane index spikes to a 6  
244 year high at the mass mortality horizon and a 12 year high in the year following the mass  
245 mortality horizon indicating a relative increase in the prevalence of water column stratification  
246 (Fig. 4g).

247

248 Thus the interval associated with the mass mortality horizon evidences decreased sulphur burial  
249 relative to carbon, higher TOC values and greater proportions of biomarkers derived from marine  
250 phytoplankton and is associated with water column stratification and hypoxic bottom waters.

251 Changes in the net burial of  $\beta,\beta$ -carotane, that might indicate a decrease in water column salinity,  
252 are coincident with the mass mortality horizon but not uniquely associated with the horizon.

253

#### 254 **4.0 DISCUSSION**

255 Geologically high concentrations of  $\beta,\beta$ -carotane (but not anomalous in the context of the section  
256 considered in this study) are reported from across the basin, particularly for localities located in  
257 palaeogeographic positions that are far from possible tributaries, indicating the prevalence of a  
258 saline or hypersaline habitat at the centre of the lake (Duncan and Hamilton 1988). From this  
259 perspective the water budget for the lake would seem to have been closed or at least heavily  
260 restricted, and at a first consideration this contradicts the relatively high rate of discharge  
261 proposed by other workers (Marshall et al., 2007), who found that riverine discharge from the  
262 lake was relatively high. The different perspectives can be reconciled by considering the  
263 seasonality of the lake; the dry seasons created a hypersaline habitat, whilst the wet seasons  
264 potentially saw large fluxes of water move through the lake. The relative duration of the two  
265 seasons would influence the net production of  $\beta,\beta$ -carotane, with less produced during a year in  
266 which the wet season predominated and riverine discharge enhanced. The changes in  $\beta,\beta$ -  
267 carotane concentration that occur several times in the studied interval suggest that the variation  
268 associated with the mass mortality horizon is not unusual and doesn't help constrain the  
269 anomalous factors at play in the genesis of the mass mortality horizon. These values represent  
270 only the routine cycling of the lake between wet and dry conditions.

271

272 Other biomarker evidence better constrains the anomalous factors that may have contributed to  
273 the formation of the mass mortality horizon. The very high  $C_{30}/C_{28}$  sterane ratio exhibits a peak

274 value at the beginning of the mass mortality horizon, but C<sub>30</sub> steranes are present in all lamina  
275 analysed albeit in trace quantities. A literal interpretation of this parameter, similar to that used  
276 for biomarkers found in oil, would suggest that the sedimentary organic matter found within the  
277 Achanarras Limestone Member predominantly derived from marine sources (Peters et al., 2007).  
278 However, as noted earlier, other geological evidence for such a strongly marine interpretation is  
279 lacking excepting the fossils of certain fish genera (such as *Coccosteus*), that are also found in  
280 similarly-aged marine successions at other localities (Trewin 1986). The proportion of 24-*n*-  
281 propylcholestane (C<sub>30</sub> steranes) varies but infrequently exceeds a single standard deviation. This  
282 can be explained by the periodic recharging of the lake with sources of C<sub>30</sub> steranes, (either  
283 phytodetritus or living organisms) during incursions of seawater, albethey some distance from  
284 the depocentre of the lake. The most likely modern analogue for such an incursion of seawater  
285 would be a storm-tide that carried non-hypersaline, marine waters into the lake or its downstream  
286 reaches.

287

288 Studies of modern day fish-kills resulting from large storm tides indicate the complexity of  
289 elucidating a definitive kill mechanism and its consequences, and generally show that the same  
290 storm will variably impact different populations in different places (Mallin et al., 2002; Schaefer  
291 et al., 2006). Van Vrancken and O'Connell (2010) described little long term change in the fish  
292 population of the downstream reaches of a small coastal tributary in Louisiana subsequent to  
293 Hurricanes Katrina and Rita, despite widespread fish-kills being evident. Conversely, at a  
294 different locality, but still within Louisiana, Perret et al., (2010) reported long term changes in  
295 fish populations that were still evident two years later. For both cases direct poisoning of fish by  
296 intrusion of saltwater itself is not considered to be the major kill-mechanism. Instead, both rapid

297 and often localised but essentially temporary hypoxia or anoxia, and widespread longer term  
298 reductions in oxygen concentration have been proposed to be the major kill-mechanisms (Mallin  
299 et al., 2002; Buck 2005). Perret et al. (2010) also considered the release of hydrogen sulphide  
300 alongside depletions in oxygen concentration as a kill-mechanism.

301  
302 The strongest evidence for atypically hypoxic conditions coincident with the mass mortality  
303 horizon is the significantly elevated values of gammacerane index (greater than 1 standard  
304 deviation) preceding the peak values in sterane parameters (both the % C<sub>28</sub> sterane and C<sub>30</sub>/C<sub>28</sub>  
305 sterane ratio). Mechanistically the link between a higher gammacerane index and “more  
306 hypoxia” or “more anoxia” is not straight forward. Foremost, it is unlikely to represent a further  
307 reduction in the oxygen concentration of a dysoxic or hypoxic water body at a single  
308 geographical point. A further reduction in oxygen concentration in a body of water that is already  
309 anoxic over a substantial depth will have little impact on net gammacerane production (see prior  
310 discussion – a stratified and anoxic water column is a cause of gammacerane production and not  
311 an input to a process governing its rate of production). Instead of representing localised changes,  
312 the changing gammacerane index likely represents the consequence of changing environmental  
313 conditions at the lakes margins, where waters that were previously oxygenated have become  
314 anoxic, thus increasing the area of the lake capable of supporting gammacerane production.

315  
316 The main mechanisms proposed by previous workers (Mallin et al., 2002; Buck 2005; Perret et  
317 al 2010) for generating hypoxic conditions during storm tides are: a) the physical mixing of deep  
318 hypoxic and sulphidic bottom water with surface waters that can cause an immediate drop in  
319 oxygen content; b) increased oxygen demand during heterotrophic activity subsequent to an algal

320 bloom triggered by an influx of nutrients – essentially a longer term phenomena; c) the  
321 entraining of anoxic but carbon-rich sediment and pore fluids within ingressing seawater and the  
322 subsequent poisoning of surface water. Mechanism a) is a rapid process that occurs during a  
323 storm, and had this been the case for the mass mortality horizon the peak gammacerane index  
324 value might be expected to have been contemporaneous with the C<sub>30</sub> sterane maxima. The  
325 maximum gammacerane index value that coincides with the C<sub>30</sub> sterane parameter maxima  
326 occurs a season latter, indicating that the environmental change was probably not instantaneous.  
327 Therefore mechanism a) is a less likely explanation for the mass mortality horizon. Mechanism  
328 c), would be expected to have left evidence in the form silt and detrital organic matter, but this is  
329 not a distinctive feature of the mass mortality horizon (although it does occur during other  
330 intervals of the Achanarras Limestone Member). Evidence for mechanism b) is thus strongest  
331 because the greatly elevated gammacerane index, that is indicative of increased hypoxia,  
332 immediately follows a peak sterane parameter value indicative of an increased contribution of  
333 phytoplankton-derived sterols (e.g. an algal bloom).

334

335 The golden alga *Prymnesium parvum* is known to produce toxins that are responsible for fish  
336 kills (Landsberg 2010). However, in the present day this alga is largely freshwater and has not  
337 been reported as a source of C<sub>30</sub> 24-*n*-propylcholesterols. Brown tide algae such as *Aureococcus*  
338 *anophagefferens*, that are tolerant of marine salinities (Doblin et al. 2004) and are reported as  
339 sources of C<sub>30</sub> 24-*n*-propylcholesterols (Giner and Boyer 1998), are generally held not to be  
340 damaging to adult fish – except when decaying algal blooms create hypoxic conditions by  
341 depleting oxygen.

342

343 The data presented in this study are from the topmost section of the Achanarras Limestone  
344 Member that has the least diverse fish assemblage (it comprises almost entirely *Dipterus*) and the  
345 lowest abundance of fish fossils (Trewin 1986). While it is tempting to try and link storm tides,  
346 seawater-incursions, reduced biodiversity and fish kills, values of the C<sub>30</sub>/C<sub>28</sub> sterane parameter  
347 are greater in the lower sections of the Achanarras Limestone Member where fish assemblages  
348 are more diverse and contain fish with the strongest marine associations. Thus the limited fish  
349 assemblage found at the top of the Achanarras Limestone Member is most likely a product of  
350 limited but highly disruptive as opposed to continuous connection to a marine environment, and  
351 indeed previous work has suggested that regular and intermittent flooding is healthy for fish  
352 stocks because it provides juvenile fish refuges on the floodplain (Mallin et al., 2002).

353

## 354 **5.0 CONCLUSION**

355 The Achanarras Limestone Member was deposited in an environment that was periodically  
356 perturbed by incursions of marine water. The extent and frequency of this perturbation is  
357 recorded in the biomarker content of individual lamina, but at the top of the fish-bearing section  
358 of the Achanarras Limestone Member incursions were short lived and lasted less than a year and  
359 were probably analogous to a modern day storm tide. When considered as a time series, a clear  
360 chronological ordering of events can be seen within biomarker data, in which an influx of  
361 seawater was followed by a period of enhanced eutrophification. Instances of storm-induced  
362 hypoxia and anoxia, as deduced from biomarker data, are most strongly associated with  
363 intermittent perturbation by incursions of seawater, and acted as a powerful environmental  
364 selection filter favoring air breathing fish such as *Dipterus*.

365



366 **ACKNOWLEDGMENTS**

367 AOW gratefully acknowledges an EPSRC Ph.D. studentship. Isotope work was financed by a  
368 NERC grant (NIGF: IP-1149-1109) and performed at SUERC, East Kilbride. The authors wish  
369 to acknowledge C. Taylor, T. Donnelly, J.Dougans and R. McGill for technical support and N.  
370 Trewin for access to the complete Achanarras Section.

371

372 **REFERENCES CITED**

373 Andrews, S.D., Trewin, N.H., Hartley, A.J., Weedon, G.P. 2010. Solar variance recorded in  
374 lacustrine deposits from the Devonian and Proterozoic of Scotland. *J. Geol. Soc.*, v.167,  
375 p.847–856.

376 Astin, T.R., 1990, The Devonian lacustrine sediments of Orkney, Scotland; implications for  
377 climate cyclicity, basin structure and maturation history. *Journal of the Geological*  
378 *Society [London]*, v.147, p.141 – 151

379 Beadle, L.C. 1981. *The inland waters of tropical Africa. An introduction to tropical liminology,*  
380 2<sup>nd</sup> edn. London: Longman.

381 Berner R.A., and Raiswell, R., 1984, C/S method for distinguishing freshwater from marine  
382 sedimentary rocks. *Geology*, v.12, p.365 – 368.

383 Bowden, S.A., Court, R.W., Milner, D., Baldwin, E.C., Lindgren, P., Crawford, I.A., Parnell, J.,  
384 Burchell, M.J. 2008. The thermal alteration by pyrolysis of the organic component of  
385 small projectiles of mudrock during capture at hypervelocity. *Journal of Analytical and*  
386 *Applied Pyrolysis*, v.82, p. 312–314.

387 Buck E.H., 2005. Hurricanes Katrina and Rita: Fishing and aquaculture industries – Damage and  
388 recovery. Congressional Research Service, Report for Congress RS22241, Washington,  
389 D.C.

390 Chen, J., Bi, Y., Zhang, J. and Li, S., 1996, Oil-source correlation in the Fulin basin, Shengli  
391 petroleum province, East China. *Organic Geochemistry*, v.24, p.931 – 940.

392 Donovan, R.N., 1993, Evaporites in the Middle Devonian of the Orcadian basin near Berriedale,  
393 Caithness. *Scottish Journal of Geology*, v.29, p.45 – 54

394 Doblin M.A., Popels L.C., Coyne K.J., Hutchins D.A., Cary S.C. and Dobbs F.C. 2014.  
395 Transport of the Harmful Bloom Alga *Aureococcus anophagefferens* by Oceangoing  
396 Ships and Coastal Boats. *Applied and Environmental Microbiology* 70, 6495 – 6500

397 Duncan, D., and Hamilton R. F. M., 1988, Palaeolimnology and organic geochemistry of the  
398 Middle Devonian in the Orcadian Basin. Geological Society [London] Special  
399 Publication, v.40, p.173-201

400 Giner, J.-L., and Boyer G. 1998. Sterols of the brown tide alga *Aureococcus anophagefferens*.  
401 *Phytochemistry* 48, 475–477.

402 Hamilton, R.F.M., and Trewin, N.H., 1988, Environmental controls on fish faunas of the Middle  
403 Devonian of the Orcadian Basin. In McMillan N.J., Embry A.F., Glass D.J. (eds)  
404 Devonian of the World. Canadian Society of Petroleum Geologists, v.14, no.3, p.589–  
405 600.

406 Huang, W.Y., and Meinschein, W. G., 1979, Sterols as ecological indicators, *Geochimica et*  
407 *Cosmochimica Acta*, v.43, no.5, p.739 – 745

408 Jiang Z.S., and Fowler, M. G., 1985, Carotenoid derived alkanes in oils from north western  
409 China, *in*: Schenck, P. A., De Leeuw, J. W. & Lijmbach, J. W. M., eds, *Advances in*  
410 *Organic Geochemistry*, Pergamon Press, Oxford.

411 Killops, S.D., and Killops, V.J., 2005, *Introduction to Organic Geochemistry*, Second Edition.  
412 Blackwell Publishing, Oxford.

413 Kodner, R.B., Pearson, A., Summons, R.E., and Knoll, A.H., 2008, Sterols in red and green  
414 algae: quantification, phylogeny, and relevance for the interpretation of geologic  
415 steranes. *Geobiology*, v.6, p.411 – 420

416 Kushwaha, S.C., Gochnauer, M.B., Kushner, D.J., Kates, M., 1974, Pigments and isoprenoid  
417 compounds in extremely and moderately halophilic bacteria. *Canadian Journal of*  
418 *Microbiology*, v.20, no.2, p.241 – 245

419 Landsberg J.H. 2002. The Effects of Harmful Algal Blooms on Aquatic Organisms, *Reviews in*  
420 *Fisheries Science*, 10:2, 113 – 390

421 Leng, M.J., and Marshall, J.D., 2004, Palaeoclimate interpretation of stable isotope data from  
422 lake sediment archives, *Quaternary Science Reviews*, v.23, p.811 – 831

423 Leventhal, J.S., 1979, The relationship between organic carbon and sulfide sulfur in recent and  
424 ancient marine and euxinic sediments. *EOS (Trans. Amer. Geophys. Union)*, v.60, p.282

425 Mallin, M.A., Posey, M.H., McIver, M.R., Parsons, D.C., Ensign, S.H. and Alphin, T.D., 2002,  
426 *Impacts and Recovery from Multiple Hurricanes in a Piedmont–Coastal Plain River*  
427 *System*. *Bioscience*, v.52, p.999 – 1010

428 Marshall, J.E.A., Rogers, D.A., and Whiteley, M.J., 1996, Marine incursions into the Orcadian  
429 Basin, Scotland. *Journal of the Geological Society [London]*, v.153, p.451 – 466

430 Marshall, J.E.A., Astin T.R., Brown J.F., Mark-Kurik E., Lazauskiene J. 2007. Recognizing the  
431 Kačák event in the Devonian terrestrial environment and its implications for  
432 understanding land-sea interactions. *Journal of the Geological Society [London]*, v.278,  
433 p.131 – 155.

434 Moldowan, J.M., 1984, C30-steranes, novel markers for marine petroleums and sedimentary  
435 rocks. *Geochimica et Cosmochimica Acta*, v.48, p.2767 – 2768

436 Othman Wilson, A., 2012. A high-resolution record of environmental and climatic change in a  
437 lacustrine sequence from the Devonian Orcadian Basin, Scotland [Ph.D. Thesis]:  
438 University of Aberdeen.

439 Perret A.J., Kaller M.D., Kelso W.E., Rutherford D.A. 2010. Effects of Hurricanes Katrina and  
440 Rita on sport fish community abundance in eastern Atchafalaya River basin, Louisiana.  
441 *N. Amer. Fish Soc.*, v.30, p. 511–517.

442 Peters, K.E., Walters, C.E., and Moldowan, J.M., 2007, *The Biomarker Guide: Volume 2,*  
443 *Biomarkers and Isotopes in Petroleum Systems and Earth History*, Cambridge University  
444 Press.

445 Rayner, D. H., 1963, The Achanarras Limestone of the Middle Old Red Sandstone, Caithness,  
446 Scotland. *Proc. Yorks. Geol. Soc.* v.34, p.117 – 138.

447 Rohmer, M., Kokke, W.C.M.C., Fenical, W., Djerassi, C. 1980. Isolation of two new c30 sterols,  
448 (24e)-24-*n*-propylidenecholesterol and 24 $\xi$ -*n*-propylcholesterol, from a cultured marine  
449 chrysophyte. *Steroids*, v. 35, p. 219-231.

450 Rønnekleiv, M., and Liaaen-Jensen, S., 1995, Bacterial carotenoids 53\* C50-carotenoids 23;  
451 carotenoids of *Haloferax volcanii* versus other halophilic bacteria. *Biochemical*  
452 *Systematics and Ecology*, v.23, no.6, p.627 – 634

453 Schaefer, J., Mickle, P., Spaeth, J., Kreiser, B.R., Adams, S.B., Matamoros, W., Zuber, B. and  
454 Vigueira, P., 2006, Effects of Hurricane Katrina on the fish fauna of the Pascagoula  
455 River Drainage. 36th Annual Mississippi Water Resources Conference, p.62 – 68

456 Sinninghe Damsté, J.S., Kenig, F., Koopmans, M.P., Köster, J., Schouten, S., Hayes, J.M. and de  
457 Leeuw, J.W., 1995, Evidence for gammacerane as an indicator of water column  
458 stratification. *Geochimica et Cosmochimica Acta*, v.59, no.9, p.1895 – 1900

459 Stephens, N.P., and Carroll, A.R., 1999, Salinity stratification in the Permian Phosphoria sea; a  
460 proposed paleoceanographic model. *Geology*, v.27, p.899 – 902

461 Ten Haven, H.L., Rohmer, M., Rullkötter, J., and Bissert, P., 1989, Tetrahymanol, the most  
462 likely precursor of gammacerane, occurs ubiquitously in marine sediments. *Geochimica et*  
463 *Cosmochimica Acta*, v.53, no.11, p.3073 – 3079

464 Trewin, N.H., 1986, Palaeoecology and sedimentology of the Achanarras fish bed of the Middle  
465 Old Red Sandstone, Scotland, *Transactions of the Royal Society of Edinburgh*, v.77,  
466 no.1, p.21 – 46.

467 Trewin, N.H., and Thirlwall, M.F., 2002, Old Red Sandstone. *in: The Geology of Scotland* 4th  
468 Edition, Trewin, N. (Ed), The Geological Society [London]

469 Van Vrancken, J. and O'Connell, M., 2010, Effects of Hurricane Katrina on freshwater fish  
470 assemblages in a small coastal tributary of Lake Pontchartrain, Louisiana. *Transactions*  
471 *of the American Fisheries Society*, v.139, p.1723 – 1732

472 Volkman J.K. 2002 Sterols in Microorganisms. *Applied Microbiol Biotechnology* 60, 495 – 506.

473

474 **FIGURE CAPTIONS**

475

476 **Fig. 1.** Map of Scotland to showing the location of the Achanarras Quarry, Caithness, the type  
477 locality for the of the Achanarras Limestone Member where samples were obtained (denoted by  
478 letter A). Distribution of Devonian deposits in Scotland is also shown.

479  
480 **Fig.2** 191 and 125  $m/z$  ion chromatograms illustrating the abundance of fossil-carotanes, hopanes  
481 and tricyclic terpanes. The y-axis of different ion-chromatograms for each sample share the same  
482 relative scale (ion counts/a.u.).  $\gamma$  =  $\gamma$ -carotane;  $\beta\beta$  =  $\beta,\beta$ -carotane;  $C_{20}$  =  $C_{20}$  13 $\beta$ (H),14 $\alpha$ (H)  
483 tricyclic terpane;  $C_{21}$  =  $C_{21}$  13 $\beta$ (H),14 $\alpha$ (H) tricyclic terpane etc.;  $C_{29}$   $\alpha\beta$  hopane =  $C_{29}$   
484 17 $\alpha$ (H),21 $\beta$ (H) 30-norhopane;  $C_{31}$   $\alpha\beta$  S hopane =  $C_{31}$  17 $\alpha$ (H),21 $\beta$ (H) (22S) hopane etc; G =  
485 gammacerane. Data are shown for the sample at the beginning of the MMH highlighted in figure  
486 4, and for 1 year after the MMH.

487  
488 **Fig. 3.** 218  $m/z$  Ion chromatogram illustrating the relative abundances of regular steranes.  $C_{27}$   
489  $\alpha\beta\beta$  R =  $C_{27}$  5 $\alpha$ ,11 $\beta$ ,14 $\beta$  (H) 20R cholestane;  $C_{27}$   $\alpha\beta\beta$  S =  $C_{27}$  5 $\alpha$ ,11 $\beta$ ,14 $\beta$  (H) 20S cholestane  
490 etc. Region of the chromatogram containing the 5 $\alpha$ ,11 $\beta$ ,14 $\beta$  (H) 20S & 20R 24- $n$ -  
491 propylcholestanes ( $C_{30}$  steranes) is shown as an inset with the y-axis at  $\times 10$  (the y-axis is ion  
492 count/a.u.).

493  
494 **Fig. 4.** Data from the 24 consecutive laminae ordinated by time assuming that 2 lamina = 1 year.  
495 (a) C/S ratio of elemental carbon to sulphur, (b) TOC (total organic carbon), (c)  $\delta^{13}C_{carb}$ , (d)  
496  $\delta^{13}C_{org}$ , (e)  $\beta,\beta$ -carotane (concentration per g of sediment), (f)  $C_{30}/C_{28}$  sterane (ratio of  $C_{30}/C_{28}$   
497 steranes), (g) %C28 sterane (percentage  $C_{28}$  sterane), (h) GI (gammacerane index =  
498 gammacerane/  $C_{31}$  14 $\alpha$ ,17 $\beta$  (H) 22S & 22 R hopanes). The MMH is shown as a grey rectangle  
499 based on the uncertainty in determining exact positions for fish beds provided in Trewin (1986).  
500 Data from a lower section of the Fish-bearing horizon of the Achanarras Limestone Member  
501 (facies 5 from Trewin 1986) are plotted as crosses at the end of the axis. Average values and  
502 standard deviations are marked in dashed lines - middle, heavier dashed line denotes arithmetic  
503 average, lighter dashed lines denote standard deviation.

504

505





Fig 1

1 year after MMH



Beginning MMH





Fig 2



a)



b)




Fig 3




Fig 4